

PMH3 - Functional Analysis

Andrew Tulloch

Contents

Chapter 1. Banach Spaces and Linear Operators	4
1. Banach Spaces	4
2. Linear Operators	6
Chapter 2. The Hahn-Banach theorem	12
1. Zorn's Lemma	13
Chapter 3. An Introduction to Hilbert Spaces	17
1. Hilbert Spaces	17
2. Projections	19
3. Orthonormal Systems	23
4. The Stone-Weierstrass Theorem	27
Chapter 4. Uniform Boundedness and the Open Mapping Theorem	35
1. The Principle of Uniform Boundedness	35
2. The Open Mapping Theorem	39
Chapter 5. Spectral Theory	44
Chapter 6. Compact Operators	53
Chapter 7. The Hilbert Space Decomposition	65
1. The Fredholm Alternative	68

Banach Spaces and Linear Operators

1. Banach Spaces

Definition 1.1 (Norm). Let X be a vector space. A norm on X is a function $\|\cdot\| : X \mapsto \mathbb{R}$ satisfying

- $\|x\| \geq 0$ with equality if and only if $x = 0$.
- $\|\alpha x\| = |\alpha|\|x\|$.
- $\|x + y\| \leq \|x\| + \|y\|$ for all $x, y \in X$.

We call the pair $(X, \|\cdot\|)$ a **normed vector space**.

Theorem 1.2 (Reverse triangle inequality). *Let X be a normed vector space. For any $x, y \in X$, we have*

$$|\|x\| - \|y\|| \leq \|x - y\|$$

Definition 1.3 (Complete space). Let X be a normed vector space. Then X is **complete** if every Cauchy sequence in X converges to some $x \in X$.

Definition 1.4 (Banach space). A **Banach space** is a complete normed vector space.

Proposition 1.5 (Convergence). *Let $(V, \|\cdot\|)$ be a normed vector space. A sequence (x_n) in V converges to $x \in V$ if given $\epsilon > 0$, there exists N such that $\|x - x_n\| < \epsilon$ whenever $n < N$.*

Lemma 1.6. *If $x_n \rightarrow x$, then $\|x_n\| \rightarrow \|x\| \in \mathbb{R}$.*

PROOF. $|\|x_n\| - \|x\|| \leq \|x - x_n\| \rightarrow 0$. □

Proposition 1.7. *Every convergent sequence is Cauchy.*

Definition 1.8 (Banach space). A complete, normed, vector space is called a **Banach space**

Proposition 1.9. $(\mathbb{K}, |\cdot|)$ is complete.

Proposition 1.10. $(\ell^p, \|\cdot\|_p)$ is a Banach space for all $1 \leq p \leq \infty$

PROOF. A general proof outline follows.

- Use completeness of \mathbb{R} to find a candidate for the limit.
- Show this limit function is in V .

- Show that $x_n \rightarrow x$ in V .

Let $x^{(n)}$ be a Cauchy sequence in ℓ^p . Since $|x_j^{(n)} - x_j^{(m)}| \leq \|x^{(n)} - x^{(m)}\|$, we know that $x_j^{(n)}$ is a Cauchy sequence in \mathbb{K} . Hence, $\lim_{n \rightarrow \infty} x_j^{(n)} := x_j$ exists, and is our limit candidate.

We now show that $\sum_{j=1}^{\infty} |x_j|^p < \infty$. We have \square

Proposition 1.11. $(\ell([a, b]), \|\cdot\|_{\infty})$ is a Banach space

Proposition 1.12. If $1 \leq p < \infty$, then $(\ell([a, b]), \|\cdot\|_p)$ is **not** a Banach space.

PROOF. Consider a sequence of functions that is equal to one on $[0, \frac{1}{2}]$, zero on $[\frac{1}{2} + \frac{1}{n}, 1]$, and linear between. This is a Cauchy sequence that does not converge to a continuous function. \square

We've seen that $(\ell([a, b]), \|\cdot\|_p)$ is not complete for $1 \leq p < \infty$.

Theorem 1.13 (Completion). Let $(V, \|\cdot\|)$ be a normed vector space over \mathbb{K} . There exists a Banach space $(V_1, \|\cdot\|_1)$ such that $(V, \|\cdot\|)$ is isometrically isomorphic to a dense subspace of $(V_1, \|\cdot\|_1)$.

Furthermore, the space $(V_1, \|\cdot\|_1)$ is unique up to isometric isomorphisms.

PROOF. Rather straightforward - construct Cauchy sequences, append limits, quotient out (as different sequences may converge to the same limit). \square

Definition 1.14. $(V_1, \|\cdot\|_1)$ is called **the completion** of $(V, \|\cdot\|)$.

Definition 1.15 (Dense). If X is a topological space and $Y \subseteq X$, then Y is **dense** in X if the closure of Y in X equals X , that is, $\bar{Y} = X$.

Alternatively, for each $x \in X$, there exists (y_n) in Y such that $y_n \rightarrow x$.

Definition 1.16 (Isomorphism of vector spaces). Two normed vector spaces $(X, \|\cdot\|_X)$ and $(Y, \|\cdot\|_Y)$ are **isometrically isomorphic** if there is a vector space isomorphism $\Psi : X \rightarrow Y$ such that

$$\|\Psi(x)\|_Y = \|x\|_X \quad \forall x \in X$$

Example 1.17. Let $\ell_0 = \{(x_i) \mid \#\{i, x_i \neq 0\} < \infty\}$. The completion of $(\ell_0, \|\cdot\|_p)$ is $(\ell^p, \|\cdot\|_p)$, because,

- ℓ_0 is a subspace of ℓ^p ,
- It is dense, since we can easily construct a sequence in ℓ_0 converging to arbitrary $x \in \ell^p$.

Example 1.18 (L^p spaces). Let μ be the Lebesgue measure on \mathbb{R} . Let

$$\mathcal{L}^p([a, b]) = \{\text{measurable } f : [a, b] \rightarrow \mathbb{K} \mid \int_a^b |f|^p d\mu < \infty\}$$

Let $\|f\|_p = \left(\int_a^b |f|^p d\mu\right)^{1/p}$. Since $\|f\|_p = 0 \iff f = 0 \text{ a.e.}$, we quotient out by the rule $f \equiv g \iff f - g = 0 \text{ a.e.}$, and then our space of equivalence classes forms a normed vector space, denoted $L^p([a, b])$.

Theorem 1.19 (Riesz-Fischer). $(L^p([a, b]), \|\cdot\|_p)$ is the completion of $(C[a, b], \|\cdot\|_p)$, and is a Banach space.

PROOF. Properties of the Lebesgue integral. □

Remark.

- Let X be any compact topological space, let $\mathcal{C}(X) = \{f : X \rightarrow \mathbb{K} \mid f \text{ is continuous}\}$, and let $\|f\|_\infty = \sup_{x \in X} \|f(x)\|$. Then $\mathcal{C}(X, \|\cdot\|_\infty)$ is Banach.
- Let X be any topological space. Then the set of all continuous and bounded functions with the supremum norm forms a Banach space.
- Let (S, \mathcal{A}, μ) be a measure space. Then we can define the \mathcal{L}^p and L^p analogously, and they are also Banach.

2. Linear Operators

Definition 2.1 (Linear operators on normed vector spaces). Let X, Y be vector spaces over \mathbb{K} . A linear operator is a function $T : X \rightarrow Y$ such that

$$\begin{aligned} T(x + y) &= T(x) + T(y) \\ T(\alpha x) &= \alpha T(x) \end{aligned}$$

for all x, y, α .

We write $\text{Hom}(X, Y) = \{T : X \rightarrow Y \mid T \text{ is linear}\}$

Definition 2.2. $T : X \rightarrow Y$ is continuous at $x \in X$ if for all $\epsilon > 0$, there exists $\delta > 0$ such that

$$\|x - y\|_X < \delta \Rightarrow \|Tx - Ty\|_Y < \epsilon$$

Definition 2.3.

$$\mathcal{L}(X, Y) = \{T : X \rightarrow Y \mid T \text{ is linear and continuous}\}$$

Remark. If $\dim(X) < \infty$ then $\text{Hom}(X, Y) = \mathcal{L}(X, Y)$. This is **not** true if X has infinite dimension.

Definition 2.4 (Bounded linear operator). Let $T : X \rightarrow Y$ be linear, then T is **bounded** if T maps bounded sets in X to bounded sets in Y . That is: for each $M > 0$ there exists $M' > 0$ such that

$$\|x\|_X \leq M \Rightarrow \|Tx\|_Y \leq M'$$

Consider the space $\mathcal{L}(X, Y)$, the set of all linear and continuous maps between two normed vector spaces X and Y .

Theorem 2.5 (Fundamental theorem of linear operators). Let $(X, \|\cdot\|_X)$ and $(Y, \|\cdot\|_Y)$ be normed vector spaces. Let $T \in \text{Hom}(X, Y)$, the set of all linear maps from X to Y . Then the following are all equivalent.

- 1) T is uniformly continuous
- 2) T is continuous
- 3) T is continuous at 0
- 4) T is bounded
- 5) There exists a constant $c > 0$ such that

$$\|Tx\|_Y \leq c\|x\|_X \quad \forall x \in X$$

PROOF. 1) \Rightarrow 2) \Rightarrow 3) is clear.

3) \Rightarrow 4). Since T is continuous at 0, given $\epsilon = 1 > 0$, there exists δ such that

$$\|Tx - T0\| \leq 1 \quad \text{whenever} \quad \|x - 0\| \leq \delta,$$

i.e. that $\|x\| \leq \delta \Rightarrow \|Tx\| \leq 1$. Let $y \in X$. The $\|\frac{\delta y}{\|y\|}\| \leq \delta$, and so $\|T\left(\frac{\delta y}{\|y\|}\right)\| \leq 1$. Hence,

$$\frac{\delta}{\|y\|} \|Ty\| \leq 1$$

and so

$$\|Ty\| \leq \frac{\|y\|}{\delta}$$

for all $y \in X$. Thus, for all $\|y\| \leq M$, we have $\|Ty\| \leq M'$, where $M' = \frac{M}{\delta}$, and so T is **bounded**.

4) \Rightarrow 5). If T is bounded, given $M = 1 > 0$, there exists $c \geq 0$ such that $\|x\| \leq 1 \Rightarrow \|Tx\| \leq c$. Then

$$\|T\left(\frac{x}{\|x\|}\right)\| \leq c$$

Hence, $\|Tx\| \leq c\|x\|$.

5) \Rightarrow 1). If 5) holds, then

$$\|Tx - Ty\| = \|T(x - y)\| \leq c\|x - y\|.$$

So if ϵ is given, taking $\delta = \frac{\epsilon}{c}$, we have

$$\|Tx - Ty\| \leq c\|x - y\| < c\frac{\epsilon}{c} = \epsilon. \quad \square$$

Corollary. If $T \in \text{Hom}(X, Y)$, then T continuous $\iff T$ bounded $\iff \|Tx\| \leq c\|x\|$ for all $x \in X$.

Definition 2.6 (Operator norm). The **operator norm** of $T \in \mathcal{L}(x, y)$, $\|T\|$ is defined by any one of the following equivalent expressions.

- (a) $\|T\| = \inf\{c > 0 \mid \|Tx\| < c\|x\|\}$.
- (b) $\|T\| = \sup_{x \neq 0} \frac{\|Tx\|}{\|x\|}$.
- (c) $\|T\| = \sup_{\|x\| \leq 1} \|Tx\|$.
- (d) $\|T\| = \sup_{\|x\|=1} \|Tx\|$.

Proposition 2.7. The operator norm is a norm on $\mathcal{L}(x, y)$.

PROOF. The following are simple to verify.

- (a) $\|T\| \geq 0$, with equality if and only if $T = 0$.
- (b) $\|\alpha T\| = |\alpha|\|T\|$.
- (c) $\|S + T\| \leq \|S\| + \|T\|$.

□

Example 2.8 (Calculating $\|T\|$). To calculate $\|T\|$, try the following.

- 1) Make sensible calculations to find c such that

$$\|Tx\| \leq c\|x\|$$

for all $x \in X$.

- 2) Find $x \in X$ such that $\|Tx\| = c\|x\|$.

Remark. Ignore !2, Q3(b), Q8 on the practice sheet, as we will be ignoring Hilbert space theory for the time being.

Definition 2.9 (Algebraic dual). Let $(X, \|\cdot\|)$ be a normed vector space over \mathbb{K} . The **algebraic dual** of X is

$$X^* = \text{Hom}(X, \mathbb{K}) = \{\varphi : X \rightarrow \mathbb{K} \mid \varphi \text{ is linear}\}.$$

Elements of X^* are called linear functionals.

Definition 2.10 (Continuous dual (just **dual**)). The **continuous dual** (just dual) of X is

$$X' = \mathcal{L}(X, \mathbb{K}) = \{\varphi : X \rightarrow \mathbb{K} \mid \varphi \text{ is linear and continuous}\}.$$

Remark. $X^* \supseteq X'$ if $\dim(X) = \infty$.

Example 2.11. Let $(\mathcal{P}([a, b]), \|\cdot\|_\infty)$ be the normed vector space of polynomials $p : [a, b] \rightarrow \mathbb{K}$.

- (a) The functional $D : \mathcal{P}([0, 1]) \rightarrow \mathbb{K}$ given by $D(p) = p'(1)$ is linear, but **not** continuous.
- (b) The functional $I : \mathcal{P}([0, 1]) \rightarrow \mathbb{K}$ given by $I(p) = \int_0^1 p(t) dt$ is linear **and** continuous.

PROOF. (a) Linearity is clear. The $p_n(t) = t^n$ for all $t \in [0, 1]$. Then $|D(p_n)| = n\|p_n\|_\infty$. So D is not continuous, as continuity implies that there exists c such that

$$\|Tx\| \leq c\|x\|.$$

- (b) Exercise: Show $\|I\| = 1$.

□

Describing the continuous dual space X' is one of the first things to do when trying to understand a normed vector space. It is generally pretty difficult to describe X' .

Proposition 2.12 (Dual of the ℓ^p space for $(1 < p < \infty)$). Let $1 < p < \infty$. Let q be the “dual” of p , defined by $\frac{1}{q} + \frac{1}{p} = 1$. Then $(\ell^p)'$ is isometrically isomorphic to ℓ^q .

Remark (Observation before proof). Let $1 \leq p < \infty$. Let $e_i = (0, 0, \dots, 1, 0, \dots)$ where 1 is in the i -th place.

1) If $x = (x_i) \in \ell^p$, then

$$x = \sum_{i=1}^{\infty} x_i e_i$$

in the sense that the partial sums converge to x .

2) If $\varphi : \ell^p \rightarrow \mathbb{K}$ is linear and continuous, then

$$\varphi(x) = \sum_{i=1}^{\infty} x_i \varphi(e_i)$$

PROOF OF OBSERVATIONS. Let $S_n = \sum_{i=1}^n x_i e_i$. Then

$$\begin{aligned} \|x - S_n\|_p^p &= \|(0, 0, \dots, x_{n+1}, x_{n+2}, \dots)\|_p^p \\ &= \sum_{i=n+1}^{\infty} |x_i|^p \\ &\rightarrow 0 \quad \text{as it is the tail of a convergent sum.} \end{aligned}$$

Write $\varphi(x)$ as

$$\begin{aligned} \varphi(x) &= \varphi(\lim_{n \rightarrow \infty} S_n) \quad (\text{continuity}) \\ &= \lim_{n \rightarrow \infty} (\varphi(S_n)) \\ &= \lim_{n \rightarrow \infty} \varphi\left(\sum_{i=1}^n x_i e_i\right) \\ &= \lim_{n \rightarrow \infty} \sum_{i=1}^n x_i \varphi(e_i) \quad (\text{linearity}) \\ &= \sum_{i=1}^{\infty} x_i \varphi(e_i) \end{aligned}$$

□

PROOF. Define a map θ by

$$\begin{aligned} \theta : \ell^q &\rightarrow (\ell^p)' \\ y &\mapsto \varphi_y \end{aligned}$$

where $\varphi_y(x) = \sum x_i y_i$ for all $x \in \ell^p$.

(1) φ_y is linear, as $\varphi_y(x + x') = \varphi_y(x) + \varphi_y(x')$ (valid as sums converge absolutely.)

(2) φ_y is continuous, as

$$|\varphi_y(x)| = \left| \sum x_i y_i \right| \leq \sum |x_i y_i| \leq \|x\|_p \|y\|_q$$

by Hölder's inequality. From the fundamental theorem of linear operators, as $|\varphi_y(x)| \leq \|x\|_p \|y\|_q$, we have that φ_y is continuous, and that

$$\|\varphi_y\| \leq \|y\|_q \quad (\star)$$

(3) θ is linear.

(4) θ is injective, as

$$\begin{aligned} \theta(y) = \theta(y') &\Rightarrow \varphi_y = \varphi_{y'} \Rightarrow \varphi_y(x) = \varphi_{y'}(x) \quad \forall x \in \ell^p \\ &\Rightarrow \varphi_y(e_i) = \varphi_{y'}(e_i) \quad \forall i \in \mathbb{N} \Rightarrow y_i = y'_i \quad \forall i \in \mathbb{N} \Rightarrow y = y' \end{aligned}$$

(5) θ is surjective. Let $\varphi \in (\ell^p)$. Let $y = (\varphi(e_1), \dots, \varphi(e_n), \dots) = (y_1, \dots, y_n, \dots)$. We now show $y \in \ell^q$.

Let $x^{(n)} \in \ell^q$ be defined by

$$x_i^{(n)} = \begin{cases} \frac{|y_i|^q}{y_i} & \text{if } i \leq n \text{ and } y_i \neq 0 \\ 0 & \text{otherwise} \end{cases}$$

Then

$$\varphi(x^{(n)}) = \sum_{i=1}^{\infty} x_i^{(n)} \varphi(e_i) = \sum_{i=1}^n |y_i|^q \quad (\dagger)$$

by Observation 2) above.

On the other hand, we know

$$\begin{aligned} \|\varphi(x^{(n)})\| &\leq \|\varphi\| \|x^{(n)}\|_p \\ &= \|\varphi\| \left(\sum_{i=1}^{\infty} |x_i^{(n)}|^p \right)^{1/p} \\ &= \|\varphi\| \left(\sum_{i=1}^n |y_i|^{(q-1)p} \right)^{1/p} \\ &= \|\varphi\| \left(\sum_{i=1}^n |y_i|^q \right)^{1/p} \quad \text{as } 1/p + 1/q = 1. \end{aligned} \quad (\star\star)$$

Now, using (\dagger) and $(\star\star)$, we have

$$\sum_{i=1}^n |y_i|^q \leq \|\varphi\| \left(\sum_{i=1}^n |y_i|^q \right)^{1/p}$$

and so we must have

$$\|y\|_q \leq \|\varphi\| \quad (\star\star\star)$$

and so $y \in \ell^q$.

We also have, by (**),

$$\|y\|_q \leq \|\varphi_y\|$$

(6) Finally, we show that θ is an isometry. By (*) and (***), we have

$$\|\theta(y)\| = \|\varphi_y\| = \|y\|_q$$

as required. □

How big is X' ? When is $X' \neq \{0\}$? Examples suggest that X' is big with a rich structure.

The Hahn-Banach theorem

The Hahn-Banach theorem is a cornerstone of functional analysis. It is all about extending linear functionals defined on a subspace to linear functionals on the whole space, while preserving certain properties of the original functional.

Definition 0.13 (Seminorm). Let X be a vector space over \mathbb{K} . A seminorm on X is a function $p : X \rightarrow \mathbb{R}$ such that

- (1) $p(x + y) \leq p(x) + p(y) \quad \forall x, y \in X$
- (2) $p(\lambda x) = |\lambda|p(x) \quad \forall x \in X, \lambda \in \mathbb{K}$

Theorem 0.14 (General Hahn-Banach). Let X be a vector space over \mathbb{K} . Let $p : X \rightarrow \mathbb{R}$ be a seminorm on X . Let $Y \subseteq X$ be a subspace of X . If $f : Y \rightarrow \mathbb{K}$ is a linear functional such that

$$|f(y)| \leq p(y) \quad \forall y \in Y$$

then there is an extension $\tilde{f} : X \rightarrow \mathbb{K}$ such that

- \tilde{f} is linear
- $\tilde{f}(y) = f(y) \quad \forall y \in Y$
- $|\tilde{f}(x)| \leq p(x) \quad \forall x \in X$

Remark. This is great.

- Y can be finite dimensional (and we know about linear functionals on finite dimensional spaces)
- If $p(x) = \|x\|$, then

$$|\tilde{f}(x)| \leq \|x\| \quad \forall x \in X$$

and so $\tilde{f} \in X'$

Corollary. Let $(X, \|\cdot\|)$ be a normed vector space over \mathbb{K} . For each $y \in X$, with $y \neq 0$, there is $\varphi \in X'$ such that

$$\varphi(y) = \|y\| \quad \text{and} \quad \|\varphi\| = 1$$

PROOF. Fix $y \neq 0$ in X . Let $Y = \{\mathbb{K}y\} = \{\lambda y | \lambda \in \mathbb{K}\}$, a one-dimensional subspace.

Define $f : Y \rightarrow \mathbb{K}$, $f(\lambda y) = \lambda\|y\|$. This is linear. Set $p(x) = \|x\|$. Then

$$|f(\lambda y)| = p(\lambda y)$$

and so by Hahn-Banach, there exists $\tilde{f} : X \rightarrow \mathbb{K}$ such that

- \tilde{f} is linear
- $\tilde{f}(\lambda y) = \lambda \tilde{f}(y) \quad \forall \lambda \in \mathbb{K}$
- $|\tilde{f}(x)| \leq \|x\| \quad \forall x \in X$

Then we have $\tilde{f} \in X'$ and $\|\tilde{f}\| = 1$ as required. \square

1. Zorn's Lemma

Theorem 1.1 (Axiom of Choice is equivalent to Zorn's Lemma). *See handout for proof that*

$$A.C. \Rightarrow Z.L.$$

Definition 1.2 (Partially ordered set). A **partially ordered set** (poset) is a set A with a relation \leq such that

- (1) $a \leq a$ for all $a \in A$,
- (2) If $a \leq b$ and $b \leq a$ then $a = b$,
- (3) If $a \leq b$ and $b \leq c$, then $a \leq c$

Definition 1.3 (Totally ordered set). A **totally ordered set** is a poset (A, \leq) such that if $a, b \in A$ then either $a \leq b$ or $b \leq a$.

Definition 1.4 (Chain). A **chain** in a poset (A, \leq) is a totally ordered subset of A .

Definition 1.5 (Upper bound). Let (A, \leq) be a poset. An **upper bound** for $B \subseteq A$ is an element $u \in A$ such that $b \leq u$ for all $b \in B$.

Definition 1.6 (Maximal element). A **maximal element** of a poset (A, \leq) is an element $m \in A$ such that $m \leq x$ implies $x = m$, that is,

$$m \leq x \Rightarrow x = m$$

Example 1.7. Let S be any set. Let $\mathcal{P}(S)$ be the power set of S (the set of all subsets of S). Define $a \leq b \iff a \subseteq b$. Maximal element is S

Theorem 1.8 (Zorn's Lemma). *Let (A, \leq) be a poset. Suppose that every chain in A has an upper bound. Then A has (at least one) maximal element.*

Example 1.9 (Application - all vector spaces have a basis).

Definition 1.10 (Linearly independent). Let X be a vector space over \mathbb{F} . We call $B \subseteq X$ **linearly independent** if

$$\lambda_1 x_1 + \cdots + \lambda_n x_n = 0 \Rightarrow \lambda_1 = \cdots = \lambda_n = 0$$

for all finite $\{x_1, \dots, x_n\} \subseteq B$.

Definition 1.11 (Span). We say $B \subseteq X$ **spans** X if each $x \in X$ can be written as

$$x = \lambda_1 x_1 + \cdots + \lambda_n x_n$$

for some $\lambda_1, \dots, \lambda_n \in \mathbb{F}$ and $\{x_1, \dots, x_n\} \subseteq B$.

Definition 1.12 (Hamel basis). A Hamel basis is a linearly independent spanning set. Equivalently, $B \subseteq X$ is a Hamel basis if and only if each $x \in X$ can be written in exactly one way as a finite linear combination of elements of B .

Theorem 1.13. *Every vector space has a Hamel basis*

PROOF. Let $L = \{\text{linearly independent subsets}\}$, with subset ordering. Let C be a chain in L . Let $u = \bigcup_{a \in C} a$. Then

- (1) $u \in L$,
- (2) u is an upper bound for C .

So Zorn's Lemma says that L has a maximal element \mathbf{b} .

Then \mathbf{b} is a Hamel basis.

- \mathbf{b} is linearly independent.
- If $\text{Span}(\mathbf{b}) \neq X$, there exists $x \in X \setminus \text{Span}(\mathbf{b})$, and $\mathbf{b}' = \mathbf{b} \cup \{x\} \in L$ is linearly independent, contradicting maximality of \mathbf{b} .

□

Remark. If $X, \|\cdot\|$ is Banach, every Hamel basis is uncountable.

Theorem 1.14 (Hahn-Banach theorem over \mathbb{R}). *Let X be a real linear space and let $p(x)$ be a seminorm on X . Let M be a real linear subspace of X and f_0 a real-valued linear functional defined on M . Let f_0 satisfy $f_0(x) \leq p(x)$ on M . Then there exists a real valued linear functional F defined on X such that*

- (i) F is an extension of f_0 , that is, $F(x) = f_0(x)$ for all $x \in M$, and
- (ii) $F(x) \leq p(x)$ on X .

PROOF. We first show that f_0 can be extended if M has codimension one. Let $x_0 \in X \setminus M$ and assume that $\text{span}(M \cup \{x_0\}) = X$. As $x_0 \notin M$ we can write $x \in X$ uniquely in the form

$$x = m + \alpha x_0$$

for $\alpha \in \mathbb{R}$. Then for every $c \in \mathbb{R}$, the map $f_c \in \text{Hom}(X, \mathbb{R})$ given by $f_c(m + \alpha x) = f_0(m) + c\alpha$ is well defined, and $f_c(m) = f_0(m)$ for all $m \in M$. We now show that we can choose $c \in \mathbb{R}$ such that $f_c(x) \leq p(x)$ for all $x \in X$. Equivalently, we must show

$$f_0(m) + c\alpha \leq p(m + \alpha x_0)$$

for all $m \in M$ and $\alpha \in \mathbb{R}$. By positive homogeneity of p and linearity of f we have

$$\begin{aligned} f_0(m/\alpha) + c &\leq p(x_0 + m/\alpha) \quad \alpha > 0 \\ f_0(-m/\alpha) - c &\leq p(-x_0 - m/\alpha) \quad \alpha < 0 \end{aligned}$$

Hence we need to choose c such that

$$\begin{aligned} c &\leq p(x_0 + m) - f_0(m) \\ c &\geq -p(-x_0 + m) + f_0(m). \end{aligned}$$

This is possible if

$$-p(-x_0 + m_1) + f_0(m_1) \leq p(x_0 + m_2) - f_0(m_2)$$

for all $m_1, m_2 \in M$. By subadditivity of p we can verify this condition since

$$f_0(m_1 + m_2) \leq p(m_1 m_2) = p(m_1 - x_0 + m_2 - x_0) \leq p(m_1 - x_0) + p(m_2 + x_0)$$

for all $m_1, m_2 \in M$. Hence c can be chosen as required.

Hence $D(F) = X$, and the theorem is proven. \square

Theorem 1.15 (Hahn-Banach over \mathbb{C}). *Suppose that c is a seminorm on a complex vector space X and let M sub a subspace of X . If $f_0 \in \text{Hom}(M, \mathbb{C})$ is such that $|f_0(x)| \leq p(x)$ for all $x \in M$, then there exists an extension $f \in \text{Hom}(X, \mathbb{C})$ such that $f|_M = f_0$ and $|f(x)| \leq p(x)$ for all $x \in X$.*

PROOF. Split f_0 into real and imaginary parts

$$f_0(x) = g_0(x) + ih_0(x).$$

By linearity of f_0 we have

$$\begin{aligned} 0 &= if_0(x) - f_0(ix) = ig_0(x) - h_0(x) - g_0(ix) - ih_0(ix) \\ &= -(g_0(ix) + h_0(x)) + i(g_0(x) - h_0(ix)) \end{aligned}$$

and so $h_0(x) = -g_0(ix)$. Therefore,

$$f_0(x) = g_0(x) - ig_0(ix)$$

for all $x \in M$. We now consider X as a vector space over \mathbb{R} , $X_{\mathbb{R}}$. Now considering $M_{\mathbb{R}}$ as a subspace of $X_{\mathbb{R}}$. Since $g_0 \in \text{Hom}(M_{\mathbb{R}}, \mathbb{R})$ and $g_0(x) \leq |f_0(x)| \leq p(x)$ and so by the real Hahn-Banach, there exists $g \in \text{Hom}(X_{\mathbb{R}}, \mathbb{R})$ such that $g|_{M_{\mathbb{R}}} = g_0$ and $g(x) \leq p(x)$ for all $x \in X_{\mathbb{R}}$. Now set $F(x) = g(x) - ig(ix)$ for all $x \in X_{\mathbb{R}}$. Then by showing $f(ix) = if(x)$, we have that f is linear.

We now show $|f(x)| \leq p(x)$. For a fixed $x \in X$ choose $\lambda \in \mathbb{C}$ such that $\lambda f(x) = |f(x)|$. Then since $|f(x)| \in \mathbb{R}$ and by definition of f , we have

$$|f(x)| = \lambda f(x) = f(\lambda x) = g(\lambda x) \leq p(\lambda x) = |\lambda| p(x) = p(x)$$

as required.

□

An Introduction to Hilbert Spaces

1. Hilbert Spaces

Definition 1.1 (Inner product). Let X be a vector space over \mathbb{K} . An **inner product** is a function

$$\langle \cdot, \cdot \rangle : X \times X \rightarrow \mathbb{K}$$

such that

- (1) $\langle x + y, z \rangle = \langle x, z \rangle + \langle y, z \rangle$
- (2) $\langle \alpha x, z \rangle = \alpha \langle x, z \rangle$
- (3) $\langle x, y \rangle = \overline{\langle y, x \rangle}$
- (4) $\langle x, x \rangle \geq 0$ with equality if and only if $x = 0$

We then have

$$\langle x, y + z \rangle = \langle x, y \rangle + \langle x, z \rangle$$

and

$$\langle x, \alpha z \rangle = \bar{\alpha} \langle x, z \rangle$$

Definition 1.2 (Inner product space). Let $(X, \langle \cdot, \cdot \rangle)$ be an **inner product space**. Defining $\|x\| = \sqrt{\langle x, x \rangle}$ turns X into a normed vector space. To prove the triangle inequality, we use the Cauchy-Swartz theorem.

Theorem 1.3 (Cauchy-Schwarz). *In an inner product space $(X, \langle \cdot, \cdot \rangle)$, we have*

$$|\langle x, y \rangle| \leq \|x\| \|y\| \quad \forall x, y \in X$$

PROOF.

$$\begin{aligned} 0 &\leq \langle x - \lambda y, x - \lambda y \rangle \\ &= \langle x, x \rangle - \langle x, \lambda y \rangle - \langle \lambda y, x \rangle + \langle \lambda y, \lambda y \rangle \\ &= \|x\|^2 - \bar{\lambda} \langle x, y \rangle - \lambda \langle y, x \rangle + |\lambda|^2 \|y\|^2 \\ &= \|x\|^2 - 2\operatorname{Re}(\lambda \langle y, x \rangle) + |\lambda|^2 \|y\|^2 \end{aligned}$$

Set $\lambda = \frac{\langle x, y \rangle}{\|y\|^2}$. Then

$$\begin{aligned} 0 &\leq \|x\|^2 - 2\operatorname{Re}\left(\frac{\langle x, y \rangle}{\|y\|^2}\right) + \frac{|\langle x, y \rangle|^2}{\|y\|^2} \\ &= \|x\|^2 - \frac{|\langle x, y \rangle|^2}{\|y\|^2} \end{aligned}$$

as required. \square

Corollary.

$$\|x + y\| \leq \|x\| + \|y\|$$

Definition 1.4 (Hilbert space). If $(X, \langle \cdot, \cdot \rangle)$ is complete with respect to $\|\cdot\|$ then it is called a **Hilbert space**.

Example 1.5. (a) ℓ^2 , where $\langle x, y \rangle = \sum_{i=1}^{\infty} x_i \bar{y}_i$.

Cauchy-Schwarz then says

$$\left| \sum_{i=1}^{\infty} x_i \bar{y}_i \right| \leq \sqrt{\sum_{i=1}^{\infty} |x_i|^2} \sqrt{\sum_{i=1}^{\infty} |y_i|^2}$$

(b) $L^2([a, b])$, where $\langle f, g \rangle = \int_a^b f(x) \overline{g(x)} dx$.

Cauchy-Swartz then says

$$\left| \int_a^b f(x) \overline{g(x)} dx \right| \leq \dots$$

Definition 1.6 (Orthogonality). Let $(X, \langle \cdot, \cdot \rangle)$ be inner product spaces. Then $x, y \in X$ are orthogonal if $\langle x, y \rangle = 0$ where $x, y \neq 0$.

Theorem 1.7. Let x_1, \dots, x_n be pairwise orthogonal elements in $(X, \langle \cdot, \cdot \rangle)$. Then

$$\left\| \sum_{i=1}^n x_i \right\|^2 = \sum_{i=1}^n \|x_i\|^2$$

Theorem 1.8 (Parallelogram identity). In $(X, \langle \cdot, \cdot \rangle)$ we have

$$\|x + y\|^2 + \|x - y\|^2 = 2(\|x\|^2 + \|y\|^2) \quad (\star)$$

for all $x, y \in X$.

Remark. If $(X, \|\cdot\|)$ is a normed vector space which satisfies parallelogram identity then X is an inner product space with inner products defined by the polarisation equation

$$\langle x, y \rangle = \begin{cases} \frac{1}{4} (\|x + y\|^2 - \|x - y\|^2) & \mathbb{K} = \mathbb{R} \\ \frac{1}{4} (\|x + y\|^2 - \|x - y\|^2 + i\|x + iy\|^2 - i\|x - iy\|^2) & \mathbb{K} = \mathbb{C} \end{cases}$$

2. Projections

Definition 2.1 (Projection). Let X be a vector space over \mathbb{K} . A subset M of X is convex if for any $x, y \in M$, then

$$tx + (1 - t)y \in M \quad \forall t \in [0, 1]$$

Theorem 2.2 (Projection). Let $(\mathcal{H}, \langle \cdot, \cdot \rangle)$ be a Hilbert space. Let $M \subseteq \mathcal{H}$ be closed and convex. Let $x \in \mathcal{H}$. Then there exists a unique point $m_x \in M$ which is closest to x , i.e.

$$\|x - m_x\| = \inf_{m \in M} \|x - m\| = d$$

PROOF. For each $k \geq 1$ choose $m_k \in M$ such that

$$d^2 \leq \|x - m_k\|^2 \leq d^2 + \frac{1}{k}$$

Each m_k exists as d is defined as the infimum over all m .

Then

$$\begin{aligned} \|m_k - m_l\|^2 &= \|(m_k - x) - (m_l - x)\|^2 \\ &= 2\|m_k - x\|^2 + 2\|m_l - x\|^2 - \|m_k + m_l - 2x\|^2 \\ &\leq 2d^2 + \frac{2}{l} + 2d^2 + \frac{2}{k} - 4\left\|\frac{m_k + m_l}{2} - x\right\|^2 \end{aligned}$$

and as $m_k/2 + m_l/2 \in M$, we have $\|\frac{m_k + m_l}{2} - x\|^2 \geq d^2$. Then

$$\|m_k - m_l\|^2 \leq 2\left(\frac{1}{k} + \frac{1}{l}\right)$$

Thus (m_k) is Cauchy. So $m_k \rightarrow m_x \in M$ as \mathcal{H} is complete and M is closed. We then have

$$\|x - m_x\| = d$$

and so now we show that m_x is unique.

Suppose that there exists $m'_x \in M$ with $\|x - m'_x\| = d$. Then by the above inequality, we have

$$\|m_x - m'_x\|^2 = 2\|m_x - x\|^2 + 2\|m'_x - x\|^2 - 4\left\|\frac{m_x + m'_x}{2} - x\right\|^2 \leq 0$$

from above. □

Definition 2.3 (Projection operator). Let $(\mathcal{H}, \langle \cdot, \cdot \rangle)$ be a Hilbert space. Let $M \subseteq \mathcal{H}$ be closed and convex. Define

$$P_M : \mathcal{H} \rightarrow \mathcal{H}$$

by $P_M(x) = m_x$ from above. This is the projection of \mathcal{H} onto M .

Definition 2.4 (Orthogonal decomposition). If $S \subseteq \mathcal{H}$, let

$$S^\perp = \{x \in \mathcal{H} \mid \langle x, y \rangle = 0 \quad \forall y \in S\}.$$

We call S^\perp the orthogonal component.

Theorem 2.5 (From previous lecture). If $M \subseteq \mathcal{H}$, then the projection of \mathcal{H} onto M is

$$\begin{aligned} P_m : \mathcal{H} &\rightarrow \mathcal{H} \\ x &\mapsto m_x \end{aligned}$$

where $m_x \in M$ is the unique element with $\|x - m_x\| = \inf_{m \in M} \|x - m\|$.

Lemma 2.6. Let $M \subseteq \mathcal{H}$ be closed subspace. Then $x - P_M x \in M^\perp$ for all $x \in \mathcal{H}$.

PROOF. Let $m \in M$. We need to show $\langle x - P_M x, m \rangle = 0$. This is clear if $m = 0$. Without loss of generality, assuming $m \neq 0$, we can assume $\|m\| = 1$. Then write

$$x - P_M x = x - (P_M x + \langle x - P_M x, m \rangle m) + \langle x - P_M x, m \rangle m.$$

Let the bracketed term be m' . Then $x - m' \perp \langle x - P_M x, m \rangle m$ because

$$\begin{aligned} \langle x - m', \langle x - P_M x, m \rangle m \rangle &= \overline{\langle x - P_M x, m \rangle} \langle x - m', m \rangle \\ &= C \langle x - P_M x - \langle x - P_M x, m \rangle m, m \rangle \\ &= C(\langle x - P_M x, m \rangle - \langle x - P_M x, m \rangle \|m\|) \\ &= 0. \end{aligned}$$

So $\|x - P_M x\|^2 = \|x - m'\|^2 + |\langle x - P_M x, m \rangle|^2$. So $\|x - P_M x\|^2 \geq \|x - P_M x\|^2 + |\langle x - P_M x, m \rangle|^2$ by definition of $P_M x$. Thus,

$$\langle x - P_M x, m \rangle = 0$$

and thus $x - P_M x \in M^\perp$. □

Theorem 2.7. The following theorem is the key fundamental result. Let $(\mathcal{H}, \langle \cdot, \cdot \rangle)$ be a Hilbert space. Let M be a closed subspace of \mathcal{H} . Then

$$\mathcal{H} = M \oplus M^\perp.$$

That is, each $x \in \mathcal{H}$ can be written in exactly one way as $x = m + m^\perp$ with $m \in M, m^\perp \in M^\perp$.

PROOF. **Existence** - Let $x = P_M x + (x - P_M x)$.

Uniqueness - Let $x = x_1 + x_1^\perp, x = x_2 + x_2^\perp$ with $x_1, x_2 \in M, x_1^\perp, x_2^\perp \in M^\perp$. Then

$$x_1 - x_2 = x_2^\perp - x_1^\perp \in M^\perp$$

Then

$$\langle x_1 - x_2, x_1 - x_2 \rangle = 0 \Rightarrow x_1 = x_2.$$

Thus $x_1^\perp = x_2^\perp$. □

Corollary. *Let $M \subseteq \mathcal{H}$ be a closed subspace. Then we have*

- (a) $P_M \in \mathcal{L}(\mathcal{H}, \mathcal{H})$.
- (b) $\|P_M\| \leq 1$.
- (c) $\text{Im} P_M = M, \text{KER } P_M = M^\perp$.
- (d) $P_M^2 = P_M$.
- (e) $P_{M^\perp} = I - P_M$.

PROOF. (c), (d), (e) exercises.

(a). Let $x, y \in H$. Write $x = x_1 + x_1^\perp$ and $y = y_1 + y_1^\perp$ with $x_1, y_1 \in M$ and $x_1^\perp, y_1^\perp \in M^\perp$. Then

$$x + y = (x_1 + y_1) + (x_1^\perp + y_1^\perp)$$

and so

$$P_M(x + y) = x_1 + y_1$$

and similarly $P_M(\alpha x) = \alpha P_M x$. We also have

$$\begin{aligned} \|x\|^2 &= \|P_M x + (x - P_M x)\|^2 \\ &= \|P_M x\|^2 + \|x - P_M x\|^2 \\ &\geq \|P_M x\|^2 \end{aligned}$$

and so $\|P_M\| \leq 1$. □

If $y \in \mathcal{H}$ is fixed, then the map

$$\begin{aligned} \varphi_y : \mathcal{H} &\rightarrow \mathbb{K} \\ x &\mapsto \langle x, y \rangle \end{aligned}$$

is in \mathcal{H}' . Linearity is clear, and continuity is proven by Cauchy-Swartz,

$$|\varphi_y(x)| = |\langle x, y \rangle| \leq \|y\| \|x\|.$$

So $\|\varphi_y\| \leq \|y\|$. Since $|\varphi_y(y)| = \|y\|^2$, we then have

$$\|\varphi_y\| = \|y\|.$$

Theorem 2.8 (Riesz Representation Theorem). *Let \mathcal{H} be a Hilbert space. The map*

$$\begin{aligned} \theta : \mathcal{H} &\rightarrow \mathcal{H}' \\ y &\mapsto \varphi_y \end{aligned}$$

is a conjugate linear bijection, and $\|\varphi_y\| = \|y\|$.

PROOF. Conjugate linearity is clear.

Injectivity

$$\varphi_y = \varphi_{y'} \Rightarrow \varphi_y(x) = \varphi_{y'}(x) \quad \forall x$$

so

$$\langle x, y = \langle x, y' \rangle = 0 \quad \Rightarrow \langle y - y', y - y' \rangle = 0$$

and so $y = y'$.

Surjectivity Let $\varphi \in H'$. We now find $y \in \mathcal{H}$ with $\varphi = \varphi_y$. If $\varphi = 0$, take $y = 0$. Suppose $\varphi \neq 0$. Then $\text{KER } \varphi \neq \mathcal{H}$. But $\text{KER } \varphi$ is a closed subspace of \mathcal{H} . So

$$H = (\text{KER } \varphi) \oplus (\text{KER } \varphi)^\perp.$$

Hence $(\text{KER } \varphi)^\perp \neq \{0\}$. Pick $z \in (\text{KER } \varphi)^\perp, z \neq 0$. For each $x \in \mathcal{H}$, the element

$$x - \frac{\varphi(x)}{\varphi(z)}z \in \text{KER } \varphi$$

Note that $\varphi(z) \neq 0$ since $z \notin \text{KER } \varphi$. Then

$$\begin{aligned} 0 &= \langle x - \frac{\varphi(x)}{\varphi(z)}z, z \rangle \\ &= \langle x, z - \frac{\varphi(x)}{\varphi(z)}z \rangle \|z\|^2 \end{aligned}$$

and so

$$\varphi(x) = \langle x, \frac{\overline{\varphi(z)}}{\|z\|^2}z \rangle \quad \forall x \in \mathcal{H},$$

and so letting $y = \frac{\overline{\varphi(z)}}{\|z\|^2}z$, we have $\varphi = \varphi_y$. □

Example 2.9. From Hahn-Banach given $y \in \mathcal{H}$ there exists $\varphi \in \mathcal{H}'$ such that

$$\|\varphi\| = 1$$

and $\varphi(y) = \|y\|$. We can be very constructive in the Hilbert case, and let

$$\varphi(x) = \langle x, \frac{y}{\|y\|} \rangle$$

Example 2.10. All continuous linear functionals on $L^2([a, b])$ are of the form

$$\varphi(f) = \int_a^b f(x)\overline{g(x)} dx$$

for some $g \in L^2([a, b])$.

Example 2.11 (Adjoint operators). Let $\mathcal{H}_1, \mathcal{H}_2$ be Hilbert spaces. Let $T \in \mathcal{L}(\mathcal{H}_1, \mathcal{H}_2)$. The **adjoint** of T is $T^* \in \mathcal{L}(\mathcal{H}_2, \mathcal{H}_1)$ given by

$$\langle Tx, y \rangle_2 = \langle x, T^*y \rangle_1$$

for all $x \in \mathcal{H}_1, y \in \mathcal{H}_2$

Exercise 2.12. Check all of the above.

Exercise 2.13. Prove $T^* = \overline{T^t}$ where T^t is the transpose.

3. Orthonormal Systems

Definition 3.1 (Orthonormal system). A subset $S \subseteq \mathcal{H}$ is an **orthonormal system** (orthonormal) if

$$\langle e, e' \rangle = \delta_{e, e'} \quad \forall e, e' \in S$$

Definition 3.2 (Complete orthonormal system or Hilbert basis). An orthonormal system S is **complete** or a **Hilbert basis** if

$$\overline{\text{span } S} = \mathcal{H}$$

Remark. By Gram-Schmidt and Zorn's Lemma, every Hilbert space has a complete orthonormal system.

Example 3.3. (1) ℓ^2 . Then

$$S = \{e_i \mid i \geq 1\}$$

is orthonormal and is complete.

(2) $L^2_{\mathbb{C}}([0, 2\pi])$. Then

$$S = \left\{ \frac{1}{\sqrt{2\pi}} e^{int} \mid n \in \mathbb{Z} \right\}$$

is orthonormal and is complete. Completeness follows from Stone-Weierstrass theorem.

(3) $L^2_{\mathbb{R}}([0, 2\pi])$. Then

$$S = \left\{ \frac{1}{\sqrt{2\pi}}, \frac{1}{\sqrt{\pi}} \cos nt, \frac{1}{\sqrt{\pi}} \sin nt \mid n \geq 1 \right\}$$

is orthonormal and is complete, again by Stone-Weierstrass.

We want to look at series $\sum_{e \in S} \dots$, which is tricky if S is not countable.

Lemma 3.4. *If $\{e_k \mid k \geq 0\}$ is orthonormal, then*

$$\sum_{k=0}^{\infty} a_k e_k$$

converges in \mathcal{H} if and only if

$$\sum_{k=0}^{\infty} |a_k|^2$$

converges in \mathbb{K} .

If either series converges, then

$$\left\| \sum_{k=0}^{\infty} a_k e_k \right\|^2 = \sum_{k=0}^{\infty} |a_k|^2$$

Note. If $x_n \rightarrow x, y_n \rightarrow y$, then

$$\langle x_n, y_n \rangle \rightarrow \langle x, y \rangle$$

PROOF. If $\sum_{k=0}^{\infty} a_k e_k$ converges to x , then

$$\begin{aligned} \langle x, x \rangle &= \lim_{n \rightarrow \infty} \left\langle \sum_{k=0}^n a_k e_k, \sum_{k=0}^n a_k e_k \right\rangle \\ &= \lim_{n \rightarrow \infty} \sum_{k=0}^n |a_k|^2 \end{aligned}$$

Conversely, if $\sum_{k=0}^{\infty} |a_k|^2$ converges, then writing $x_n = \sum_{k=0}^n a_k e_k$, we have

$$\begin{aligned} \|x_m - x_n\|^2 &= \left\| \sum_{k=n+1}^m a_k e_k \right\|^2 \\ &= \sum_{k=n+1}^m \|a_k e_k\|^2 \text{ by Pythagoras} \\ &= \sum_{k=n+1}^m |a_k|^2 \rightarrow 0 \end{aligned}$$

and so (x_n) is Cauchy, and hence converges by completeness of \mathcal{H} . □

Lemma 3.5. Let $\{e_1, \dots, e_n\}$ be orthonormal. Then

$$\sum_{k=1}^n |\langle x, e_k \rangle|^2 \leq \|x\|^2$$

for each $x \in \mathcal{H}$.

PROOF. Let $y = \sum_{k=1}^n \langle x, e_k \rangle e_k$. Let $z = x - y$. We claim that $z \perp y$. We have

$$\begin{aligned} \langle x, y \rangle &= \langle x - y, y \rangle \\ &= \langle x, y \rangle - \|y\|^2 \\ &= \sum_{k=1}^n \overline{\langle x, e_k \rangle} \langle x, e_k \rangle - \sum_{k=1}^n |\langle x, e_k \rangle|^2 \\ &= 0. \end{aligned}$$

So

$$\begin{aligned}\|x\|^2 &= \|y + z\|^2 \\ &= \|y\|^2 + \|z\|^2 \text{ Pythagoras} \\ &\geq \|y\|^2 = \sum_{k=1}^n |\langle x, e_k \rangle|^2\end{aligned}$$

□

We want to write expressions like $\sum_{e \in S} \langle x, e \rangle e$.

Corollary. *Let $x \in \mathcal{H}$ and S orthonormal. Then*

$$\{e \in S \mid \langle x, e \rangle \neq 0\}$$

is countable.

PROOF.

$$\{e \in S \mid \langle x, e \rangle \neq 0\} = \bigcup_{k \geq 1} \{e \in S \mid |\langle x, e \rangle| > \frac{1}{k}\}$$

From the lemma,

$$\#\{e \in S \mid |\langle x, e \rangle| > \frac{1}{k}\} \leq k^2 \|x\|^2$$

For if this number were greater than $k^2 \|x\|^2$, then the LHS in Lemma is greater than $\frac{1}{k^2} k^2 \|x\|^2$. □

Therefore:

Corollary (Bessel's Inequality). *If S is orthonormal, then*

$$\sum_{e \in S} |\langle x, e \rangle|^2 \leq \|x\|^2$$

for all $x \in \mathcal{H}$

PROOF. $\sum_{e \in S} |\langle x, e \rangle|^2$ is a sum of countably many positive terms, and so order is not important. □

We want to write $\sum_{e \in S} \langle x, e \rangle e$. This sum is over a countable set, but is the order important?

Theorem 3.6. *Let S be orthonormal. Let $M = \overline{\text{span } S}$. Then*

$$P_M x = \sum_{e \in S} \langle x, e \rangle e$$

where the sum can be taken in any order.

PROOF. Fix $x \in H$. Choose an enumeration

$$\{e_k \mid k \geq 0\} = \{e \in S \mid \langle x, e \rangle \neq 0\}.$$

By Bessel's inequality, we have

$$\sum_{k=0}^{\infty} |\langle x, e_k \rangle|^2 \leq \|x\|^2$$

and so the LHS converges. By Lemma 3.4, we know

$$y = \sum_{k=0}^{\infty} \langle x, e_k \rangle e_k \in M$$

converges in \mathcal{H} .

Write $x = y + (x - y) = M + M^\perp$. We claim $(x - y) \in M^\perp$. Then $P_M x = y$ from characterisation of projection operator. Let $e \in S$. Then

$$\begin{aligned} \langle x - y, e \rangle &= \lim_{n \rightarrow \infty} \left\langle x - \sum_{k=0}^n \langle x, e_k \rangle e_k, e \right\rangle \\ &= \lim_{n \rightarrow \infty} (\langle x, e \rangle - \sum_{k=0}^n \langle x, e_k \rangle \langle e_k, e \rangle) \\ &= \langle x, e \rangle - \sum_{k=0}^{\infty} \langle x, e_k \rangle \langle e_k, e \rangle. \end{aligned}$$

If $e \in \{e' \in S \mid \langle x, e' \rangle \neq 0\}$, then $e = e_j$ for some j , and so

$$\langle x - y, e \rangle = \langle x, e_j \rangle - \langle x, e_j \rangle = 0$$

If $\langle x, e \rangle = 0$, then $e \neq e_j$ for all j , and so $\langle e_j, e \rangle = 0$, and so

$$\langle x - y, e \rangle = 0 - 0 = 0.$$

Thus $x - y \in (\text{span } S)^\perp$.

Exercise 3.7. Show that

$$x - y \in \overline{(\text{span } S)}^\perp = M^\perp.$$

□

Recall that if $\{x_1, \dots\}$ is a countable orthonormal system in a Hilbert space \mathcal{H} . Then

$$\sum_{k=1}^{\infty} a_k e_k < \infty \iff \sum_{k=1}^{\infty} |a_k|^2 < \infty$$

and

$$\left\| \sum_{k=1}^{\infty} a_k e_k \right\|^2 = \sum_{k=1}^{\infty} |a_k|^2 \quad (\star)$$

We also had the following.

Theorem 3.8. Let S be orthonormal in \mathcal{H} . Let $M = \overline{\text{span } S}$. Then

$$P_M x = \sum_{e \in S} \langle x, e \rangle e \quad \forall x \in \mathcal{H}$$

where the sum has only countable many terms and convergence is unconditional.

Theorem 3.9. Let S be orthonormal in \mathcal{H} . Then following are equivalent.

- (a) S is a complete orthonormal system ($\overline{\text{span } S} = \mathcal{H}$).
- (b) $x = \sum_{e \in S} \langle x, e \rangle e$ for all x (Fourier series).
- (c) $\|x\|^2 = \sum_{e \in S} |\langle x, e \rangle|^2$ for all x (Parseval's formula).

PROOF. (a) \Rightarrow (b). If $M = \overline{\text{span } S} = \mathcal{H}$, then

$$P_M x = x = \sum_{e \in S} \langle x, e \rangle e$$

by Theorem 3.8.

(b) \Rightarrow (c). By the infinite Pythagoras theorem (\star).

(c) \Rightarrow (a). Let $M = \overline{\text{span } S}$. Suppose that $z \in M^\perp$. Then $z = 0 + z \in M + M^\perp$. Hence

$$0 = \|P_M z\|^2 = \left\| \sum_{e \in S} \langle z, e \rangle e \right\|^2 = \sum_{e \in S} |\langle z, e \rangle|^2 = \|z\|^2$$

which implies $z = 0$, so $M = \mathcal{H}$, and so S is complete. \square

Remark. Consider $L^2([0, 2\pi])$, and let $S = \{e_n \mid n \in \mathbb{Z}\}$. Then we can write

$$f = \sum_{n \in \mathbb{Z}} c_n e_n$$

where $c_n = \langle f, e_n \rangle = \frac{1}{\sqrt{2\pi}} \int_0^{2\pi} f(t) e^{-int} dt$.

We do not claim that convergence is pointwise, what we have proven is convergence in L^2 ,

$$\left\| f - \sum_{|n| \leq N} c_n e_n \right\|_2 \rightarrow 0$$

as $N \rightarrow \infty$. This is not the same as pointwise or uniform convergence ($\|\cdot\|_\infty$).

4. The Stone-Weierstrass Theorem

This is a useful tool to show an orthonormal system is complete. In fact, this theorem is about uniformly approximating elements of $\mathcal{C}(X)$, where X is a compact Hausdorff space. It is a generalisation of the Weierstrass approximation theorem.

Theorem 4.1 (Weierstrass approximation theorem). Let $f \in \mathcal{C}([a, b])$ and let $\epsilon > 0$ be given. Then there exists a polynomial $p(x)$ such that

$$|f(x) - p(x)| < \epsilon \quad \forall x \in [a, b],$$

that is, $\|f - p\|_\infty < \epsilon$.

Corollary. *This implies the following important results:*

- *Continuous functions can be uniformly approximated by polynomials.*
- $\mathcal{P}([a, b])$, *the space of polynomials on $[a, b]$, is dense in $\mathcal{C}([a, b])$.*
- $\overline{\mathcal{P}([a, b])} = \mathcal{C}([a, b])$.

We now prove Stone's 1930's generalisation.

First some setup: Let X be a compact Hausdorff space throughout. We then know that $\mathcal{C}(X)$ is a vector space. It also has sensible vector multiplication,

$$(fg)(x) = f(x)g(x).$$

Thus $\mathcal{C}(X)$ is a unital, commutative, associative ring. As we have

$$f(\lambda g) = \lambda(fg)$$

then $\mathcal{C}(X)$ is a unital, commutative, associative algebra over \mathbb{K} .

Definition 4.2 (Subalgebra). A subalgebra of $\mathcal{C}(X)$ is a subset \mathcal{A} which is closed under scalar multiplication, vector addition, and vector multiplication. \mathcal{A} is unital if it contains the constant function $f(x) = 1$.

Example 4.3. $\mathcal{P}([a, b])$ is a subalgebra of $\mathcal{C}([a, b])$.

When is \mathcal{A} dense in $\mathcal{C}(X)$?

Theorem 4.4 (Stone-Weierstrass theorem). *Let X be a compact Hausdorff space, and let \mathcal{A} be a subalgebra of $\mathcal{C}(X)$. If*

- (1) \mathcal{A} *is unital,*
- (2) $f \in \mathcal{A} \Rightarrow f^* \in \mathcal{A}$, *where $f^*(x) = \overline{f(x)}$,*
- (3) \mathcal{A} *separates points of X .*

Then $\overline{\mathcal{A}} = \mathcal{C}(X)$.

Definition 4.5. \mathcal{A} separates points of X if, given $x \neq y$, there is a function $f \in \mathcal{A}$ with $f(x) \neq f(y)$.

Corollary. (a) $\mathcal{P}([a, b])$ *is dense in $\mathcal{C}([a, b])$, as $f(x) = x$ separates points.*

(b) *Trigonometric polynomials are dense in*

$$\{f \in \mathcal{C}([0, 2\pi]) \mid f(0) = f(2\pi)\}.$$

(c) *Trigonometric polynomials are dense in $L^2([0, 2\pi])$, and*

$$S = \{e_n \mid n \in \mathbb{Z}\}$$

is complete.

Setup

Lemma 4.6. *The function $f(t) = |t|$ can be uniformly approximated by polynomials on $[-1, 1]$*

PROOF. The binomial theorem says

$$(1+x)^{1/2} = \sum_{n=0}^{\infty} \binom{1/2}{n} x^n \quad \forall x \in [-1, 1]$$

We then have

$$|t| = \sqrt{t^2} = \sqrt{1 + (t^2 - 1)} = \sum_{n=0}^{\infty} \binom{1/2}{n} (t^2 - 1)^n \quad t \in [-\sqrt{2}, \sqrt{2}]$$

Now let $p_N(t) = \sum_{n=0}^N \binom{1/2}{n} (t^2 - 1)^n$, and

$$||t| - p_N(t)| = \left| \sum_{n=N+1}^{\infty} \binom{1/2}{n} (t^2 - 1)^n \right| \leq \sum_{n=N+1}^{\infty} \left| \binom{1/2}{n} \right|$$

and so $||t| - p_N|_{\infty} \rightarrow 0$ as $N \rightarrow \infty$ on $[-1, 1]$. \square

Theorem 4.7 (Stone-Weierstrass theorem). *Let X be a compact Hausdorff space, and let \mathcal{A} be a subalgebra of $\mathcal{C}(X)$. If*

- (1) \mathcal{A} is unital,
- (2) $f \in \mathcal{A} \Rightarrow f^* \in \mathcal{A}$, where $f^*(x) = \overline{f(x)}$,
- (3) \mathcal{A} separates points of X .

Then $\overline{\mathcal{A}} = \mathcal{C}(X)$.

PROOF. We first prove for $\mathcal{C}_{\mathbb{R}}(X)$.

Lemma 4.8. *Let \mathcal{A} be a unital subalgebra of $\mathcal{C}_{\mathbb{R}}(X)$. Then*

- (a) $|f| \in \overline{\mathcal{A}}$,
- (b) $\min(f_1, \dots, f_n), \max(f_1, \dots, f_n) \in \overline{\mathcal{A}}$

for all $f, f_1, \dots, f_n \in \mathcal{A} \subseteq \mathcal{C}_{\mathbb{R}}(X)$.

PROOF. (a) Replace f by $\frac{f}{\|f\|_{\infty}}$ so we can assume that $\|f\|_{\infty} = 1$. From the previous lemma, we know for each $n \geq 1$ there is a polynomial $p_n : [-1, 1] \rightarrow \mathbb{R}$ such that $||t| - p_n(t)| < \frac{1}{n}$ for all $t \in [-1, 1]$.

Since $|f(x)| \leq \|f\|_{\infty} = 1$ for all $x \in X$, we have

$$||f| - p_n(f)| \leq \frac{1}{n}$$

But $p_n(f)$ is a finite linear combination of $1, f, f^2, f^3, \dots$ and so in \mathcal{A} , as \mathcal{A} is unital. Thus $|f| \in \overline{\mathcal{A}}$.

(b) Use the formulas

$$\max(f, g) = \frac{f + g + |f - g|}{2}, \quad \min(f, g) = \frac{f + g - |f - g|}{2} \in \overline{\mathcal{A}}$$

and induction. □

PROOF OF STONE-WEIERSTRASS FOR $\mathcal{C}_{\mathbb{R}}(X)$. Let $f \in \mathcal{C}_{\mathbb{R}}(X)$ and let $\epsilon > 0$ be given. We need to find $g \in \mathcal{A}$ such that

$$|f(z) - g(z)| < \epsilon \quad \forall z \in X$$

Step 0. We can assume that \mathcal{A} is closed.

Exercise 4.9. Why?

Step 1. Let $x, y \in X$ be fixed.

Proposition 4.10. *There exists $f_{xy} \in \mathcal{A}$ with*

$$f_{xy}(x) = f(x), \quad f_{xy}(y) = f(y)$$

PROOF. If $x = y$ then trivial (take $f_{xy}(z) = f(x)\mathbf{1}(z)$).

If $x \neq y$, since \mathcal{A} separates points, there is $h \in \mathcal{A}$ with $h(x) \neq h(y)$. Then take

$$f_{xy} = ah + b\mathbf{1} \in \mathcal{A}$$

we can invert the coefficient matrix to find our coefficients a and b . □

Step 2. Let $x \in X$ be fixed.

Proposition 4.11. *There exists $f_x \in \mathcal{A}$ such that*

- $f_x(x) = f(x)$.
- $f_x(z) < f(z) + \epsilon$

PROOF. For each $y \in X$, let

$$O_y = \{z \in X \mid f_{xy}(z) < f(z) + \epsilon\}$$

where f_{xy} is the function from Step 1. These are all open sets (why?) and thus

$$X = \bigcup_{y \in X} O_y$$

since $x \in O_y$.

By compactness of X , we have

$$X = \bigcup_{i=1}^m O_{y_i}$$

Letting $f_x = \min(f_{xy_1}, \dots, f_{xy_m})$. Then

- Since $f_{x_{y_i}}(x) = f(x)$ for all i ,

$$f_x(x) = f(x)$$

- If $z \in X$, then $z \in O_{y_i}$ for some i , and so

$$f_x(z) \leq f_{x_{y_i}}(z) < f(z) + \epsilon$$

as required. □

Step 3.

Proposition 4.12. *There exists a function $g \in \mathcal{A}$ such that*

$$|f(z) - g(z)| < \epsilon$$

for all $z \in X$.

PROOF. For each $x \in X$, let

$$U_x = \{z \in X \mid f_x(z) > f(x) - \epsilon\}$$

where f_x is from Step 2. These sets U_x are open and since $x \in U_x$, for an open cover, we can write

$$X = \bigcup_{x \in X} U_x = \bigcup_{j=1}^n U_{x_j}.$$

Define $g = \max(f_{x_1}, \dots, f_{x_n})$. If $z \in X$,

- $g(z) = f_{x_i}(z)$ for some i , which is less than $f(z) + \epsilon$ from Step 2.
- If $z \in U_{x_j}$ for some $j = 1, \dots, n$, then

$$g(z) \geq f_{x_j}(z) > f(x) - \epsilon.$$

□

□

Exercise 4.13. Where did we use the Hausdorff property?

We now prove for $\mathcal{C}_{\mathbb{C}}(X)$.

Let

$$\mathcal{A}_{\mathbb{R}} = \{f \in \mathcal{A} \mid f \text{ is real valued}\}.$$

Then $\mathcal{A}_{\mathbb{R}}$ is an \mathbb{R} -subalgebra of $\mathcal{C}_{\mathbb{R}}(X)$. It is unital, as $1 \in \mathcal{A}$ and it is real valued.

We now show $\mathcal{A}_{\mathbb{R}}$ separates points. If $x \neq y$, there is $f \in \mathcal{A}$ such that $f(x) \neq f(y)$. Write $f = u + iv$ with u, v real valued. Either $u(x) \neq u(y)$ or $v(x) \neq v(y)$, and so $\mathcal{A}_{\mathbb{R}}$ separates points.

Hence $\mathcal{A}_{\mathbb{R}}$ is dense in $\mathcal{C}_{\mathbb{R}}$.

Now, let $f \in \mathcal{C}_{\mathbb{C}}(X)$. Then write $f = u + iv$. Then $u, v \in \mathcal{C}_{\mathbb{R}}(X)$. Then given $\epsilon > 0$, there exists $u_1, v_1 \in \mathcal{A}_{\mathbb{R}}$ such that

$$\|u - u_1\|_{\infty} \leq \frac{\epsilon}{2}, \quad \|v - v_1\|_{\infty} \leq \frac{\epsilon}{2}$$

Writing $f_1 = u_1 + iv_1 \in \mathcal{A}$, we have

$$\|f - f_1\|_{\infty} \leq \|(u - u_1) + i(v - v_1)\|_{\infty} \leq \|u - u_1\|_{\infty} + \|v - v_1\|_{\infty} < \epsilon$$

and thus \mathcal{A} is dense in $\mathcal{C}_{\mathbb{C}}(X)$. \square

Corollary. *Polynomials are dense in $\mathcal{C}([a, b])$.*

PROOF. $\mathcal{A} = \mathcal{P}([a, b])$ is an algebra, is unital, is closed under complex conjugation, and separates points. Thus, \mathcal{A} is dense in $\mathcal{C}([a, b])$. \square

Definition 4.14 (Trigonometric polynomials). A **trigonometric polynomial** is an expression

$$\sum_{n \in \mathbb{Z}} c_n e^{int}$$

with finitely many $c_n \neq 0$. So these are polynomials in $s = e^{it}$ and $s^{-1} = \bar{s} = e^{-it}$.

Corollary. *The space \mathcal{A} of all trigonometric polynomials is dense in $\mathcal{C}(\Pi)$, where $\Pi = \{z \in \mathbb{C} \mid |z| = 1\}$*

PROOF. \mathcal{A} is a sub-algebra of $\mathcal{C}(\Pi)$, it is unital, closed under complex conjugation,

$$\overline{\sum_{n \in \mathbb{Z}} c_n e^{int}} = \sum_{n \in \mathbb{Z}} \overline{c_{-n}} e^{int}$$

and separates points. T is a compact Hausdorff space, and thus Stone-Weierstrass states that \mathcal{A} is dense in $\mathcal{C}(\Pi)$. \square

Corollary. *The orthonormal system*

$$S = \left\{ \frac{1}{\sqrt{2\pi}} e^{int} \mid n \in \mathbb{Z} \right\}$$

is complete in $L^2([0, 2\pi])$.

PROOF. $\text{span } S = \mathcal{A}$ is a space of trigonometric polynomials, which is dense in $\mathcal{C}(\Pi)$. Define

$$\begin{aligned} \Phi : \mathcal{C}_p([0, 2\pi]) &\rightarrow \mathcal{C}(\Pi) \\ f &\mapsto \tilde{f} \end{aligned}$$

where $\mathcal{C}_p([0, 2\pi]) = \{f \in \mathcal{C}([0, 2\pi]) \mid f(0) = f(2\pi)\}$. Then Φ is an isometric isomorphism, and therefore functions of the form $f(t) = \sum c_n e^{int}$ is dense in $\mathcal{C}_p([0, 2\pi])$.

By the construction of the Lebesgue integral, **simple** functions

$$\sum_{i=1}^n a_i \mathbf{1}_{A_i}$$

are dense in $L^2([0, 2\pi])$.

Exercise 4.15. Given $f \in L^2([0, 2\pi])$ and $\epsilon > 0$, there exists $g \in \mathcal{C}_p([0, 2\pi])$ such that $\|f - g\|_2 < \epsilon$.

Thus \mathcal{A} is dense in $L^2([0, 2\pi])$. □

Corollary. *The following are separable (have a countable dense subset):*

- (a) $\mathcal{C}([a, b])$,
- (b) $L^p([a, b])$, $1 \leq p < \infty$

PROOF. (a) We have $\mathcal{P}([a, b])$ is dense in $\mathcal{C}([a, b])$ and set $\mathcal{P}_{\mathbb{Q}}([a, b])$ with rational coefficients is dense in $\mathcal{P}([a, b])$. Clearly, $\mathcal{P}_{\mathbb{Q}}([a, b])$ is countable, and thus is dense in $\mathcal{C}([a, b])$.

(b) Use the fact that $\mathcal{C}([a, b])$ is dense in $L^p([a, b])$. □

Corollary. *Let X be a compact metric space. Then $\mathcal{C}(X)$ is separable.*

PROOF. As X is a compact metric space, then X is separable.

Exercise 4.16. Why?

Let $\{x_n \mid n \geq 1\}$ be a countable dense subset of X . For each $n \geq 1$ and $m \geq 1$ define

$$f_{n,m} : X \rightarrow \mathbb{K}$$

by

$$f_{n,m}(x) = \inf_{z \notin B(x_n, \frac{1}{m})} d(x, z)$$

We then claim $f_{n,m}$ is continuous. Now, let \mathcal{A} be the space of all \mathbb{K} -linear combinations of

$$f_{n_1, m_1}^{k_1}, \dots, f_{n_l, m_l}^{k_l}, k_1, \dots, k_l \in \mathcal{N}. \quad (\star)$$

This is a sub-algebra of $\mathcal{C}(X)$, as \mathcal{A} is unital, closed under conjugation, and separates points - if $z_1, z_2 \in X$ with $z_1 \neq z_2$, Choose n, m such that $z_1 \in B(x_n, \frac{1}{m})$, $z_2 \notin B(x_n, \frac{1}{m})$. Thus the sub-algebra \mathcal{A} is dense by Stone-Weierstrass.

The subset of \mathbb{Q} -linear combinations of (\star) is countable and dense. □

Lemma 4.17. *If X is compact metric space then X is separable.*

PROOF. For each $m \geq 1$,

$$X = \bigcup_{x \in X} B(x; \frac{1}{m})$$

has a finite subcover

$$X = \bigcup_{n=1}^{N_m} B(x_{m,n}, \frac{1}{m})$$

and thus the subset of all $\{x_{m,n}\}$ is a countably dense subset. \square

Corollary.

$$\frac{\pi^2}{6} = \sum_{n=1}^{\infty} \frac{1}{n^2}.$$

PROOF. $S = \{\frac{1}{\sqrt{2\pi}}e^{int} \mid n \in \mathbb{Z}\}$ is complete, and so Parseval's formula holds,

$$\|f\|_2^2 = \sum_{n \in \mathbb{Z}} |\langle f, e_n \rangle|^2.$$

Apply to $f(x) = x$. \square

A common strategy is to prove for polynomials, and then Stone-Weierstrass proves it for continuous functions.

Corollary. *If $f \in \mathcal{C}([a, b] \times [c, d])$ then*

$$\int_a^b \int_c^d f(x, y) dy dx = \int_c^d \int_a^b f(x, y) dx dy$$

PROOF. By direct calculation, the result is true for two-variable polynomials. Let $f \in \mathcal{C}([a, b] \times [c, d])$ and $\epsilon > 0$ be given. By Stone-Weierstrass, the space of polynomials in 2 variables is dense in $\mathcal{C}([a, b] \times [c, d])$ and so there exists a polynomial $p(x, y)$ with

$$|f(x, y) - p(x, y)| < \frac{\epsilon}{(b-a)(d-c)}.$$

The result then follows by direct calculation. \square

Uniform Boundedness and the Open Mapping Theorem

The following is at the core of two of the cornerstone theorems of functional analysis - the uniform boundedness principle and the open mapping theorem.

1. The Principle of Uniform Boundedness

Theorem 1.1 (Baire's theorem). *Let X be a complete metric space. If U_1, U_2, \dots are open dense subsets of X , then*

$$U = \bigcap_{n=1}^{\infty} U_n$$

is dense in X .

PROOF. Let $x \in X$ and $\epsilon > 0$ be given. We need to show that

$$B(x, \epsilon) \cap U \neq \emptyset.$$

Lemma 1.2. *There exists sequences (x_n) in X and (ϵ_n) in \mathbb{R}^+ with the property that*

- (a) $x_1 = x, \epsilon_1 = \epsilon.$
- (b) $\epsilon_n \downarrow 0$
- (c) $\overline{B(x_{n+1}, \epsilon_{n+1})} \subseteq B(x_n, \epsilon_n) \cap U_n$ for all $n \geq 1.$

PROOF. Let x_1, \dots, x_n and $\epsilon_1, \dots, \epsilon_n$ be chosen. By density of U_n ,

$$B(x_n, \epsilon_n) \cap U_n \neq \emptyset.$$

Choose $x_{n+1} \in B(x_n, \epsilon_n) \cap U_n$. Choose $\epsilon'_{n+1} > 0$ such that $B(x_{n+1}, \epsilon'_{n+1}) \subseteq B(x_n, \epsilon_n) \cap U_n$ (openness). We have $\epsilon'_{n+1} \leq \epsilon_n$. Choose $0 < \epsilon_{n+1} \leq \min(\frac{\epsilon'_{n+1}}{2}, \frac{1}{n+1})$, then we have

$$\begin{aligned} \overline{B(x_{n+1}, \epsilon_{n+1})} &\subseteq B(x_{n+1}, \epsilon'_{n+1}) \\ &\subseteq B(x_n, \epsilon_n) \cap U_n \end{aligned}$$

and $\epsilon_{n+1} < \epsilon_n$ with $\epsilon_{n+1} < \frac{1}{n+1}$. □

Given the lemma, the theorem follows. If $m \geq n$, then by (c),

$$B(x_m, \epsilon_m) \subseteq B(x_n, \epsilon_n) \cap U_n \tag{*}$$

In particular, $x_m \in B(x_n, \epsilon_n)$. Thus, $d(x_n, x_m) < \epsilon_n$ for all $m \geq n$. Thus (x_n) is Cauchy, and so $x_n \rightarrow \zeta$ in X by completeness. By (\star) , we then have $d(x_n, \zeta) \leq \epsilon_n$ for all $n \geq 1$. So $\zeta \in \overline{B(x_n, \epsilon_n)}$. So by (c), $\zeta \in \overline{B(x_{n+1}, \epsilon_{n+1})} \subseteq B(x_n, \epsilon_n) \cap U_n$.

Thus $\zeta \in B(x, \epsilon)$ and thus $\zeta \in U = \bigcap_{n=1}^{\infty} U_n$. \square

The following corollary is often used

Corollary. *Let X be a complete metric space. If C_1, C_2, \dots are closed with $X = \bigcup_{n=1}^{\infty} C_n$ then $\text{Int}(C_n) \neq \emptyset$ for some n .*

PROOF. If $\text{Int}(C_n) = \emptyset$ for all n then $U_n = X \setminus C_n$ are open and dense. So by Baire's theorem, $\bigcap_{n=1}^{\infty} U_n$ is dense, and in particular, $\bigcap_{n=1}^{\infty} U_n \neq \emptyset$. We have

$$\begin{aligned} X &= \bigcup_{n=1}^{\infty} C_n = \bigcup_{n=1}^{\infty} (X \setminus U_n) \\ &= X \setminus \left(\bigcap_{n=1}^{\infty} U_n \right) \\ &\subsetneq X, \end{aligned}$$

a contradiction. \square

There are three cornerstone theorems.

- Hahn-Banach,
- Uniform Boundedness,
- Open Mapping.

Theorem 1.3 (Uniform boundedness). *Let X, Y be Banach spaces. Let $T_\alpha, \alpha \in A$, a family of continuous linear operators $T_\alpha : X \rightarrow Y$. Then if*

$$\sup_{\alpha \in A} \|T_\alpha x\| < \infty$$

for each fixed $x \in X$, then

$$\sup_{\alpha \in A} \|T_\alpha\| < \infty$$

Remark. Rather amazing - you get a global bound from pointwise bounds.

PROOF. For each $n \geq 1$, let

$$X_n = \{x \in X \mid \|T_\alpha x\| \leq n \forall \alpha \in A\}$$

These are **closed** (T_α is continuous) and

$$X = \bigcup_{n=1}^{\infty} X_n$$

by the hypothesis.

By the corollary to Baire's theorem, we know there exists $n_0 \geq 1$ with $\text{Int}(X_{n_0}) \neq \emptyset$. Choose $x_0 \in \text{Int}(X_{n_0})$, and let $r > 0$ such that

$$B(x_0, r) \subseteq \text{Int}(X_{n_0}).$$

If $\|z\| \leq 1$ then $x_0 + rz \in \overline{B}(x_0, r)$. So $x_0 + rz \in X_{n_0}$, and

$$\|T_\alpha(x_0 + rz)\| \leq n_0 \quad \forall \alpha \in A,$$

but $\|a\| - \|b\| \leq \|a + b\|$, so

$$\|T_\alpha(rz)\| - \|T_\alpha(x_0)\| \leq \|T_\alpha(x_0 + rz)\| \leq n_0.$$

So $r\|T_\alpha z\| \leq n_0 + n_0$, and

$$\|T_\alpha z\| \leq \frac{2n_0}{r} \quad \forall \|z\| \leq 1, \forall \alpha \in A$$

For a general $x \in X$,

$$\|T_\alpha x\| = \|T_\alpha\left(\frac{x}{\|x\|}\right)\| \|x\| \leq \frac{2n_0}{r} \|x\|$$

and thus $\|T_\alpha\| \leq \frac{2n_0}{r}$, which implies

$$\sup_{\alpha \in A} \|T_\alpha\| < \infty \quad \square$$

Recall, the Fourier series of $f \in L^2([-\pi, \pi])$ is

$$\sum_{k \in \mathbb{Z}} \langle f, e_k \rangle e_k$$

where $e_k(t) = \frac{e^{ikt}}{\sqrt{2\pi}}$. This converges to f in the L^2 norm.

Exercise 1.4. If f is 2π -periodic and continuous, does the Fourier series converge pointwise?

There are explicit (complicated) examples, but the easiest existence is using the uniform boundedness principle.

Proposition 1.5. *There is a 2π periodic continuous function whose Fourier series does not converge at 0.*

PROOF. Let $\mathcal{C}_p([-\pi, \pi]) = \{f \in \mathcal{C}([-\pi, \pi]) \mid f(-\pi) = f(\pi)\}$. This is a Banach space with $\|\cdot\|_\infty$. If $f \in \mathcal{C}_p$, let

$$f_n = \sum_{|k| \leq n} \langle f, e_k \rangle e_k.$$

Remark. We can now define, for each $n \geq 1$, a linear operator $T_n : \mathcal{C}_p \rightarrow \mathbb{K}$ by

$$T_n(f) = f_n(0).$$

If $f_n(0)$ converges (as $n \rightarrow \infty$) for each $f \in \mathcal{C}_p$, then

$$\sup_{n \geq 1} |T_n f| = \sup_{n \geq 1} |f_n(0)| < \infty$$

for all $f \in \mathcal{C}_p$, which by uniform boundedness implies

$$\sup_{n \geq 1} \|T_n\| \leq \infty. \quad (\star)$$

We now show that (\star) is false.

We have

$$\begin{aligned} f_n(x) &= \sum_{|k| \leq n} \frac{1}{2\pi} \int_{-\pi}^{\pi} f(t) e^{-ikt} dt e^{ikx} \\ &= \frac{1}{2\pi} \int_{-\pi}^{\pi} f(t) \left(\sum_{|k| \leq n} e^{-ik(x-t)} \right) dt \\ &= \frac{1}{2\pi} \int_{-\pi}^{\pi} f(t) D_n(x-t) dt \end{aligned}$$

where $D_n(t) = \sum_{|k| \leq n} e^{ikt}$ is the **Dirichlet Kernel**. The Dirichlet kernel is real, and even, with

$$D_n(t) = \frac{\sin(n + \frac{1}{2})t}{\sin \frac{t}{2}}.$$

Note. T_n is continuous, with norm $\|T_n\| = \frac{1}{2\pi} \int_{-\pi}^{\pi} |D_n(t)| dt$.

PROOF.

$$\begin{aligned} |T_n(f)| &\leq \frac{1}{2\pi} \int_{-\pi}^{\pi} |f(t)| |D_n(t)| dt \\ &\leq \left(\frac{1}{2\pi} \int_{-\pi}^{\pi} |D_n(t)| dt \right) \|f\|_{\infty} \end{aligned}$$

and so $\|T_n\| \leq \frac{1}{2\pi} \int_{-\pi}^{\pi} |D_n(t)| dt$.

Going the other way, let

$$s_t = \begin{cases} 1 & D_n(t) \geq 0 \\ -1 & D_n(t) < 0 \end{cases}.$$

We have seen that set functions can be approximated in L^1 -norm by continuous (periodic) functions.

So if $\epsilon > 0$ is given, there is a $g \in \mathcal{C}_p$ such that

$$\left| \frac{1}{2\pi} \int_{-\pi}^{\pi} (g(t) - s(t)) D_n(t) dt \right| < \epsilon.$$

g can be chosen with $\|g\|_{\infty} = 1$.

So

$$\left| T_n(g) - \frac{1}{2\pi} \int_{-\pi}^{\pi} |D_n(t)| dt \right| < \epsilon.$$

Thus

$$\frac{1}{2\pi} \int_{-\pi}^{\pi} |D_n(t)| dt - |T_n(g)| < \epsilon.$$

So

$$|T_n(g)| \geq \frac{\|g\|_{\infty}}{2\pi} \int_{-\pi}^{\pi} |D_n(t)| dt - \epsilon.$$

Since $\epsilon > 0$ was arbitrary,

$$\|T_n\| \geq \frac{1}{2\pi} \int_{-\pi}^{\pi} |D_n(t)| dt. \quad \square$$

All that remains is to show that

$$\|T_n\| = \frac{1}{2\pi} \int_{-\pi}^{\pi} |D_n(t)| dt \rightarrow \infty$$

We have

$$\begin{aligned} \|T_n\| &= \frac{1}{\pi} \int_0^{\pi} |D_n(t)| dt \\ &= \frac{1}{\pi} \int_0^{\pi} \frac{|\sin(n + \frac{1}{2})t|}{|\sin \frac{t}{2}|} dt \\ &\geq \frac{2}{\pi} \int_0^{\pi} \frac{|\sin(n + \frac{1}{2})t|}{t} dt \\ &= \frac{2}{\pi} \int_0^{(n+\frac{1}{2})\pi} \frac{\sin v}{v} dv \\ &\geq \frac{2}{\pi} \int_0^{n\pi} \frac{\sin v}{v} dv \\ &= \frac{2}{\pi} \sum_{k=1}^n \int_{(k-1)\pi}^{k\pi} \frac{|\sin v|}{v} dv \\ &\geq \sum_{k=1}^n \frac{1}{k\pi} \int_{(k-1)\pi}^{k\pi} |\sin v| dv \\ &= \frac{4}{\pi^2} \sum_{k=1}^n \frac{1}{k} \rightarrow \infty \end{aligned}$$

as $n \rightarrow \infty$.

Thus there exists $f \in \mathcal{C}_p$ such that the Fourier series of f diverges at $x = 0$. □

2. The Open Mapping Theorem

This theorem is *tailor-made* to deal with inverse operators.

Definition 2.1 (Open mapping). Let X, Y be metric spaces. A function $f : X \rightarrow Y$ is **open** if open sets in X are mapped to open sets in Y .

Theorem 2.2 (Open mapping theorem). Let X, Y be Banach spaces. If $T \in \mathcal{L}(X, Y)$ is surjective then T is open.

Corollary (Bounded inverse theorem). Let X, Y be Banach spaces. If $T \in \mathcal{L}(X, Y)$ is bijective, then

$$T^{-1} \in \mathcal{L}(Y, X).$$

PROOF. Let $O \subseteq X$ be open. Then $(T^{-1})^{-1}(O) = T(O)$ is open (by the open mapping theorem). Thus T^{-1} is continuous. \square

Corollary. Let $(X, \|\cdot\|_1)$ and $(X, \|\cdot\|_2)$ be Banach spaces. If

$$\|x\|_1 \leq C\|x\|_2 \quad \forall x \in X$$

then $\|\cdot\|_1$ and $\|\cdot\|_2$ are equivalent.

PROOF.

$$\begin{aligned} i : (X, \|\cdot\|_2) &\rightarrow (X, \|\cdot\|_1) \\ x &\mapsto x \end{aligned}$$

is linear, surjective and injective, and also continuous, as

$$\|i(x)\| = \|x\|_1 \leq C\|x\|_2.$$

So the bounded inverse theorem gives

$$i^{-1} : (X, \|\cdot\|_1) \rightarrow (X, \|\cdot\|_2)$$

is continuous. Thus there exists $A > 0$ such that $\|i^{-1}(x)\|_2 \leq A\|x\|_1$, which implies $\|x\|_2 \leq A\|x\|_1$. So

$$\frac{1}{A}\|x\|_2 \leq \|x\|_1 \quad \forall x \in X \quad \square$$

More generally, if $T \in \mathcal{L}(X, Y)$ is bijective, then by the bounded inverse theorem,

$$c\|x\| \leq \|Tx\| \leq C\|x\|$$

where $c = \frac{1}{\|T^{-1}\|}$, $C = \|T\|$.

Lemma 2.3. Let X be a Banach space and Y a normed space. Then for $T \in \mathcal{L}(X, Y)$, the following are equivalent.

- (a) T is open
- (b) There exists $r > 0$ such that $B(0, r) \subseteq T(\overline{B(0, 1)})$

(c) There exists $r > 0$ such that $B(0, r) \subseteq \overline{T(\overline{B(0, 1)})}$.

PROOF. (a) \Rightarrow (b), (c). As $B(0, 1)$ is open, the set $T(B(0, 1))$ is open in Y . Since $0 \in T(B(0, 1))$ there exists $\delta > 0$ such that the set

$$B(0, \delta) \subseteq T(B(0, 1)) \subseteq \overline{T(\overline{B(0, 1)})} \subseteq \overline{T(\overline{B(0, 1)})}.$$

(c) \Rightarrow (b). Assume that there exists $r > 0$ such that

$$B(0, r) \subseteq \overline{T(\overline{B(0, 1)})}.$$

We now show that $B(0, \frac{r}{2}) \subseteq \overline{T(\overline{B(0, 1)})}$ which proves (b). Let $y \in B(0, \frac{r}{2})$. Then $2y \in B(0, r)$ and since $B(0, r) \subseteq \overline{T(\overline{B(0, 1)})}$ there exists $x_1 \in \overline{B(0, 1)}$ such that

$$\|2y - Tx_1\| \leq \frac{r}{2}$$

Hence $4y - 2Tx_1 \in B(0, r)$ and by the same argument as before there exists $x_2 \in \overline{B(0, 1)}$ such that

$$\|4y - 2Tx_1 - Tx_2\| \leq \frac{r}{2}$$

Continuing this way we construct a sequence $(x_n) \in \overline{B(0, 1)}$ such that

$$\|2^n y - 2^{n-1}Tx_1 - \dots - 2Tx_{n-1} - Tx_n\| \leq \frac{r}{2}$$

for all n . Dividing by 2^n we obtain

$$\|y - \sum_{k=1}^n 2^{-k}Tx_k\| \leq \frac{r}{2^{n+1}}$$

Hence $y = \sum_{k=1}^{\infty} 2^{-k}Tx_k$. Since $\|x_k\| \leq 1$ for all $k \in \mathbb{N}$ we have that

$$\sum_{k=1}^{\infty} 2^{-k}\|x_k\| \leq \sum_{k=1}^{\infty} 2^{-k} = 1$$

and so the series

$$x = \sum_{k=1}^{\infty} 2^{-k}x_k$$

converges absolutely in X as X is Banach and hence complete. We have also that $\|x\| \leq 1$ and so $x \in \overline{B(0, 1)}$. Because T is continuous we have

$$Tx = \lim_{n \rightarrow \infty} \sum_{k=1}^n 2^{-k}Tx_k = y$$

by construction of x . Hence $y \in \overline{T(\overline{B(0, 1)})}$ and (b) follows.

(b) \Rightarrow (a). By (b) and the linearity of T we have

$$T(\overline{B(0, \epsilon)}) = \epsilon \overline{T(\overline{B(0, 1)})}$$

for all $\epsilon > 0$. Since the map $x \mapsto \epsilon x$ is a homeomorphism on Y the set $T(\overline{B(0, \epsilon)})$ is a neighbourhood of zero for all $\epsilon > 0$. Now let $U \subseteq X$ be open and $y \in T(U)$. As U is open there exists $\epsilon > 0$ such that

$$\overline{B(x, \epsilon)} = x + \overline{B(0, \epsilon)} \subseteq U$$

where $y = Tx$. Since $z \mapsto x + z$ is a homeomorphism and T is linear we have

$$T(\overline{B(x, \epsilon)}) = Tx + T(\overline{B(0, \epsilon)}) = y + T(\overline{B(0, \epsilon)}) \subseteq T(U).$$

Hence $T(\overline{B(x, \epsilon)})$ is a neighbourhood of y in $T(U)$. As y was arbitrary in $T(U)$ it follows that $T(U)$ is open. \square

Lemma 2.4. *Let X be a normed vector space and $S \subseteq X$ convex with $S = -S$. If \overline{S} has a non-empty interior, then \overline{S} is a neighbourhood of zero.*

PROOF. First note that \overline{S} is convex. If $x, y \in S$ and $x_n, y_n \in S$ with $x_n, y_n \rightarrow x, y$ then $tx_n + (1 - ty_n) \in S$ for all n and $t \in [0, 1]$. Letting $n \rightarrow \infty$ we get $tx + (1 - t)y \in \overline{S}$ for all $t \in [0, 1]$ and so \overline{S} is convex. We also easily have $\overline{S} = -\overline{S}$. If \overline{S} has a non-empty interior, there exists $z \in \overline{S}$ and $\epsilon > 0$ such that $B(z, \epsilon) \subseteq \overline{S}$. Therefore $z \pm h \in \overline{S}$ whenever $\|h\| < \epsilon$ and since $\overline{S} = -\overline{S}$ we also have $-(z \pm h) \in \overline{S}$. By the convexity of \overline{S} we have

$$y = \frac{1}{2}((x + h) + (-x + h)) \in \overline{S}$$

whenever $\|h\| < \epsilon$. Hence $B(0, \epsilon) \subseteq \overline{S}$, and so \overline{S} is a neighbourhood of zero. \square

Theorem 2.5 (Open mapping theorem). *Suppose that X and Y are Banach spaces. If $T \in \mathcal{L}(X, Y)$ is surjective, then T is open.*

PROOF. As T is surjective we have

$$Y = \bigcup_{n \in \mathbb{N}} \overline{T(\overline{B(0, n)})}$$

with $\overline{T(\overline{B(0, n)})}$ closed for all $n \in \mathbb{N}$. Since Y is complete, by a corollary to Baire's theorem, there exists $n \in \mathbb{N}$ such that $\overline{T(\overline{B(0, n)})}$ has non-empty interior. Since the map $x \mapsto nx$ is a homeomorphism and T is linear, the set $\overline{T(\overline{B(0, 1)})}$ has non-empty interior as well. Now $\overline{B(0, 1)}$ is convex and $\overline{B(0, 1)} = -\overline{B(0, 1)}$. By linearity of T we have that

$$\overline{T(\overline{B(0, 1)})} = -\overline{T(\overline{B(0, 1)})}$$

is convex as well. Since we know that $\overline{T(\overline{B(0, 1)})}$ has non-empty interior, the previous lemma implies that $\overline{T(\overline{B(0, 1)})}$ is a neighbourhood of zero, and thus there exists $r > 0$ such that

$$B(0, r) \subseteq \overline{T(\overline{B(0, 1)})}$$

and since X is Banach the previous lemma shows that T is open. \square

Exercise 2.6. If X, Y are vector spaces, and if $T : X \rightarrow Y$ is linear, then $\Gamma(T)$ is a subspace of $X \times Y$. Moreover, if X, Y are normed vector spaces, with

$$\|(x, Tx)\|_{\Gamma} = \|x\| + \|Tx\|.$$

Theorem 2.7 (Closed Graph theorem). *Let X, Y be Banach spaces, and $T \in \text{Hom}(X, Y)$. Then $T \in \mathcal{L}(X, Y)$ if and only if $\Gamma(T)$ is closed in $X \times Y$.*

PROOF. Suppose $T \in \mathcal{L}(X, Y)$. If $x_n \rightarrow x$ in X , then

$$(x_n, Tx_n) \rightarrow (x, Tx)$$

by continuity of T , and so $\Gamma(T)$ is closed.

Conversely, suppose that $\Gamma(T)$ is closed in $X \times Y$. Define a norm $\|\cdot\|_{\Gamma}$ on X by $\|x\|_{\Gamma} = \|x\| + \|Tx\|$. Since $\Gamma(T)$ is closed, and since $(X, \|\cdot\|)$ is Banach, then $(X, \|\cdot\|_{\Gamma})$ is also a Banach space (exercise). Note that $\|x\| \leq \|x\|_{\Gamma}$. So by a corollary to the Open Mapping theorem, $\|\cdot\|$ and $\|\cdot\|_{\Gamma}$ are equivalent. So there is $c > 0$ with

$$\|x\|_{\Gamma} \leq c\|x\| \quad \forall x \in X.$$

So $\|x\| + \|Tx\| \leq c\|x\|$, and so $\|Tx\| \leq (c - 1)\|x\|$, and so T is continuous. \square

Spectral Theory

The eigenvalues of an $n \times n$ matrix T over \mathbb{C} are the $\lambda \in \mathbb{C}$ with

$$\det(\lambda I - T) = 0$$

that is, $\lambda I - T$ is not invertible.

Remark. Showing existence of eigenvalues is equivalent to the fundamental theorem of algebra.

Remark. We need our base field to be \mathbb{C} to get reasonable spectral theory.

Definition 0.8. Write $\mathcal{L}(X) = \mathcal{L}(X, X)$.

Definition 0.9. Let X be a Banach space over \mathbb{K} , and let $T \in \mathcal{L}(X)$. Then the spectrum of T is

$$\sigma(T) = \{\lambda \in \mathbb{K} \mid \lambda I - T \text{ is not invertible}\}.$$

Remark. $\lambda I - T$ is non invertible if either $\lambda I - T$ is not injective, or $\lambda I - T$ is not surjective.

Remark. If $\dim(X) < \infty$, then $X \setminus \text{KER}(T) \simeq \text{im}(T)$, and so T is injective if and only if T is surjective. This fails in the infinite dimensional case - consider the left and right shift operators on ℓ^2 .

Definition 0.10 (Eigenvalue). $\lambda \in \mathbb{K}$ is an eigenvalue of $T \in \mathcal{L}(X)$ if there is $x \neq 0$ with $Tx = \lambda x$, i.e. λ is an eigenvalue if and only if $\lambda I - T$ is not injective.

Theorem 0.11. Let $X \neq \{0\}$ be a Banach space over \mathbb{C} , and let $T \in \mathcal{L}(X)$. Then $\sigma(T)$ is a non-empty, compact (closed and bounded) subset of

$$\{\lambda \in \mathbb{C} \mid |\lambda| \leq \|T\|\}$$

Example 0.12. Let $L, R : \ell^2 \rightarrow \ell^2$ be the left and right shift operators.

Then $\|L\| = 1$, and so $\sigma(L) \subseteq \overline{D}(0, 1)$. If $|\lambda| < 1$, then

$$L(\lambda, \lambda^2, \lambda^3, \dots) = (\lambda^2, \lambda^3, \lambda^4, \dots) = \lambda(\lambda, \lambda^2, \lambda^3, \dots)$$

and so λ is an eigenvalue. Thus $D(0, 1) \subseteq \sigma(L) \subseteq \overline{D}(0, 1)$. But $\sigma(L)$ is closed, and so $\sigma(L) = \overline{D}(0, 1)$. Are the λ with $|\lambda| = 1$ eigenvalues? No - suppose $|\lambda| = 1$ and $x \neq 0$ with $Lx = \lambda x$.

Then

$$L^n(x) = \lambda^n x.$$

Thus, $x_{n+1} = \lambda^n x_1$. Then $x = (x_1, \lambda x_1, \lambda^2 x_1, \dots)$ which is not in ℓ^2 .

Then $\|R\| = 1$, and so $\sigma(R) \subseteq \overline{D}(0, 1)$.

Note. $LRx = L(0, x_1, \dots) = (x_1, x_2, \dots)$, so

$$LR = I \tag{*}$$

Remark. Unlike $\dim(X) < \infty$, (*) does NOT say that R is invertible ($RL = I$).

Consider the operator $L(\lambda I - R) = \lambda L - I = -\lambda(\lambda^{-1}I - L)$. If $0 < |\lambda| < 1$, then we know that $\lambda^{-1}I - L$ is invertible (as $\lambda^{-1} \notin \sigma(L)$). So if $\lambda I - R$ were invertible, then L is invertible, which is false. Thus $\lambda \in \sigma(R)$. Hence

$$D(0, 1) \setminus \{0\} \subseteq \sigma(R) \subseteq \overline{D}(0, 1).$$

Since $\sigma(R)$ is closed, $\sigma(R) = \overline{D}(0, 1)$.

Theorem 0.13. *Let $X \neq \{0\}$ be a Banach space over \mathbb{C} . Let $T \in \mathcal{L}(X)$. Then $\sigma(T)$ is a nonempty, compact subset of*

$$\{\lambda \in \mathbb{C} \mid |\lambda| \leq \|T\|\}.$$

Lemma 0.14. *With above assumptions $\sigma(T) \subseteq \{\lambda \in \mathbb{C} \mid |\lambda| \leq \|T\|\}$.*

PROOF. We need to show that if $|\lambda| > \|T\|$ then $\lambda I - T$ is invertible.

Technique: Geometric series. We guess

$$(\lambda I - T)^{-1} = \frac{1}{\lambda I - T} = \sum_{k=0}^{\infty} \frac{T^k}{\lambda^{k+1}}.$$

We now verify this guess. Since

$$\sum_{k=0}^{\infty} \frac{\|T^k\|}{|\lambda|^{k+1}} \leq \sum_{k=0}^{\infty} \frac{\|T\|^k}{|\lambda|^{k+1}} < \infty,$$

the series $S = \sum_{k=0}^{\infty} \frac{T^k}{\lambda^{k+1}}$ converges in X .

We now show that S is the inverse of $\lambda I - T$. As we are working in infinite dimensions, we need to check left and right inverses. Let $S_n = \sum_{k=0}^{n-1} \frac{T^k}{\lambda^{k+1}}$. Then

$$\begin{aligned} S_n(\lambda I - T) &= \left(\sum_{k=0}^{n-1} \frac{T^k}{\lambda^{k+1}} \right) (\lambda I - T) \\ &= I - \frac{T^n}{\lambda^n} \rightarrow I \\ (\lambda I - T)S_n &= I - \frac{T^n}{\lambda^n} \rightarrow I \end{aligned}$$

and so $S(\lambda I - T) = (\lambda I - T)S$ and so $\lambda I - T$ is invertible. \square

Exercise 0.15. Show that if $\|I - T\| < 1$ then T is invertible with inverse $\sum_{k=0}^{\infty} (I - T)^k$ *Hint: Consider*

$$\frac{1}{T} = \frac{1}{I - (I - T)}.$$

In particular, the ball $B(I, 1)$ in $\mathcal{L}(X)$ consists of invertible elements.

The following is used to show $\sigma(T)$ is closed and nonempty, it is also interesting in its own right.

Proposition 0.16. *Let X be Banach over \mathbb{K} . Let $GL(X) = \{T \in \mathcal{L}(X) \mid T \text{ invertible}\}$. Then*

- (a) $GL(X)$ is a group under composition of operators.
- (b) $GL(X)$ is open in $\mathcal{L}(X)$.
- (c) The map

$$\begin{aligned} \varphi : GL(X) &\rightarrow GL(X) \\ T &\mapsto T^{-1} \end{aligned}$$

is continuous.

PROOF. (a) The open mapping theorem tells us that if $T \in GL(X)$ then $T^{-1} \in \mathcal{L}(X)$, and so $T^{-1} \in GL(X)$. The rest is clear.

- (b) Let $T_0 \in GL(X)$. We claim

$$B\left(T_0, \frac{1}{\|T_0^{-1}\|}\right) \subseteq GL(X).$$

We have

$$\begin{aligned} \|I - T_0^{-1}T\| &= \|T_0^{-1}(T_0 - T)\| \\ &\leq \|T_0^{-1}\| \|T_0 - T\| \\ &< 1 \quad \text{as } T \in B\left(T_0, \frac{1}{\|T_0^{-1}\|}\right) \end{aligned}$$

- (c) We have

$$\begin{aligned} \|T_0^{-1} - T^{-1}\| &= \|T^{-1}(T - T_0)T_0^{-1}\| \\ &\leq \|T^{-1}\| \|T - T_0\| \|T_0^{-1}\| \end{aligned} \tag{*}$$

If $\|T - T_0\| \leq \frac{1}{2\|T_0^{-1}\|}$, then

$$\begin{aligned}\|I - TT_0^{-1}\| &= \|(T_0 - T)T_0^{-1}\| \\ &\leq \|T_0 - T\|\|T_0^{-1}\| \\ &\leq \frac{1}{2}.\end{aligned}$$

We then have

$$\begin{aligned}\|T_0T^{-1}\| &= \|(TT_0^{-1})^{-1}\| \\ &= \left\| \sum_{k=0}^{\infty} (I - TT_0^{-1})^k \right\| \\ &\leq \sum_{k=0}^{\infty} \|I - TT_0^{-1}\|^k \\ &\leq 2\end{aligned}$$

Hence $\|T^{-1}\| = \|T_0^{-1}(T_0T^{-1})\| \leq \|T_0^{-1}\|\|T_0T^{-1}\| \leq 2\|T_0^{-1}\|$, and from (\star) , we have

$$\|T_0^{-1} - T^{-1}\| \leq 2\|T_0^{-1}\|^2\|T - T_0\|$$

and so $T \mapsto T^{-1}$ is continuous. □

Corollary. $\sigma(T)$ is closed.

PROOF. Let

$$\begin{aligned}f: \mathbb{C} &\rightarrow \mathcal{L}(X) \\ \lambda &\mapsto \lambda I - T\end{aligned}$$

This is continuous, as

$$\begin{aligned}\|f(\lambda) - f(\lambda_0)\| &= \|(\lambda - \lambda_0)I\| \\ &= |\lambda - \lambda_0|\end{aligned}$$

and

$$\sigma(T) = f^{-1}(\mathcal{L}(X) \setminus \text{GL}(X))$$

which is the inverse image of a closed set, and hence is closed. □

So $\sigma(T)$ is a compact subset of $\{\lambda \in \mathbb{C} \mid |\lambda| \leq \|T\|\}$. Write $\rho(T) = \mathbb{C} \setminus \sigma(T)$ (the **resolvent set**), and let $R_T = R: \rho(T) \rightarrow \mathcal{L}(X)$ with $R_T(\lambda) = (\lambda I - T)^{-1}$.

Theorem 0.17. Let $\mathbb{K} = \mathbb{C}$ and $X \neq \{0\}$ and $T \in \mathcal{L}(X)$. Then $\sigma(T) \neq \emptyset$.

PROOF. We use Liouville's theorem - a bounded entire function must be constant.

Let $\varphi = \mathcal{L}(X)'$ (hence $\varphi : \mathcal{L}(X) \rightarrow C$.) Let

$$\begin{aligned} f_\varphi &: \rho(T) \rightarrow \mathbb{C} \\ \lambda &\mapsto \varphi(R(\lambda)) \end{aligned}$$

Lemma 0.18. f_φ is analytic on $\rho(T)$.

PROOF. We show f_φ is differentiable. Consider

$$\begin{aligned} \frac{f_\varphi(\lambda) - f_\varphi(\lambda_0)}{\lambda - \lambda_0} &= \varphi \left(\frac{R(\lambda) - R(\lambda_0)}{\lambda - \lambda_0} \right) \\ &= \varphi \left(\frac{(\lambda I - T)^{-1} - (\lambda_0 I - T)^{-1}}{\lambda - \lambda_0} \right) \\ &= \varphi \left(\frac{(\lambda_0 I - T)^{-1}((\lambda_0 - \lambda)I)(\lambda I - T)^{-1}}{\lambda - \lambda_0} \right) \\ &= -\varphi \left((\lambda_0 I - T)^{-1}(\lambda I - T)^{-1} \right) \\ &\rightarrow -\varphi \left((\lambda_0 I - T)^{-2} \right) \end{aligned}$$

as $\lambda \rightarrow \lambda_0$, where we use the fact that φ is continuous and $T \rightarrow T^{-1}$ is continuous. So f_φ is analytic on $\rho(T)$ for all $\varphi \in \mathcal{L}(X)'$. \square

Now suppose that $\sigma(T) = \emptyset$. Then $f_\varphi : \mathbb{C} \rightarrow \mathbb{C}$ is analytic.

Lemma 0.19. f_φ is bounded.

PROOF. If $|\lambda| > \|T\|$, then

$$\begin{aligned} f_\varphi(\lambda) &= \left| \varphi \left((\lambda I - T)^{-1} \right) \right| \\ &= \left| \varphi \left(\sum_{k=0}^{\infty} \frac{T^k}{\lambda^{k+1}} \right) \right| \\ &\leq \|\varphi\| \left\| \sum_{k=0}^{\infty} \frac{T^k}{\lambda^{k+1}} \right\| \\ &\leq \|\varphi\| \sum_{k=0}^{\infty} \frac{\|T\|^k}{|\lambda|^{k+1}} \\ &= \frac{\|\varphi\|}{|\lambda| - \|T\|} \rightarrow 0 \end{aligned}$$

as $|\lambda| \rightarrow \infty$. So f_φ is bounded, entire, and thus $f_\varphi = c$ by Liouville's theorem. By the above, $f_\varphi(\lambda) = 0$ for all λ . Hence $\varphi(R(\lambda)) = 0$ for all λ, φ .

Thus from Hahn-Banach, $R(\lambda) = 0$ for all λ which is a contradiction, as the zero operator is not invertible if $X \neq \{0\}$. \square

□

Theorem 0.20 (Spectral mapping theorem (polynomials)). *Let T be an $n \times n$ matrix over \mathbb{C} . If we know all the eigenvalues of T , then we know the eigenvalues of every polynomial $p(T) = a_0 + a_1T + \cdots + a_nT^n$. Specifically,*

$$\{\text{eigenvalues of } p(T)\} = \{p(\lambda) \mid \lambda \text{ is an eigenvalue of } T\}$$

Therefore

$$\sigma(p(T)) = p(\sigma(T)).$$

This is called the **spectral mapping theorem** (for matrices/polynomials).

This also holds for X Banach over \mathbb{C} , and $T \in \mathcal{L}(X)$.

Lemma 0.21. *Let $\mathbb{C}[t]$ be the algebra of polynomials in t with complex coefficients. Multiplication is defined as usual.*

Lemma 0.22. *Let X be Banach over \mathbb{C} . Let $T \in \mathcal{L}(X)$. Then*

$$\begin{aligned} \varphi : \mathbb{C}[t] &\rightarrow \mathcal{L}(X) \\ p &\mapsto p(T) \end{aligned}$$

is an algebra homomorphism (multiplication corresponds to composition in $\mathcal{L}(X)$.)

PROOF. Simply check

$$\begin{aligned} \varphi(p_1 + p_2) &= \varphi(p_1) + \varphi(p_2) \\ \varphi(p_1 p_2) &= \varphi(p_1) \varphi(p_2) \\ \varphi(\alpha p) &= \alpha \varphi(p) \end{aligned}$$

for all $p_1, p_2, p \in \mathbb{C}[t], \alpha \in \mathbb{C}$. □

Theorem 0.23. *Let X be Banach over \mathbb{C} , and let $T \in \mathcal{L}(X)$. Then*

$$\sigma(p(T)) = p(\sigma(T)).$$

PROOF. If $p = c$ is constant, then $p(T) = cI$ has spectrum

$$\sigma(p(T)) = \sigma(cI) = \{c\}$$

On the other hand,

$$p(\sigma(T)) = \{c\}$$

Now, suppose that p is non constant. Let $\mu \in \mathbb{C}$ fixed. By the fundamental theorem of algebra, we can factorise $\mu - p(t)$ as

$$\alpha(t - \lambda_1)^{m_1} \cdots (t - \lambda_n)^{m_n}$$

where $\lambda_1, \dots, \lambda_n$ are the distinct roots of $\mu - p(t)$. Note that $\mu = p(\lambda_i)$ for each i . Applying $\psi : \mathbb{C}[t] \rightarrow \mathcal{L}(X)$ from above, we have

$$\mu I - p(T) = \alpha(T - \lambda_1 I)^{m_1} \dots (T - \lambda_n I)^{m_n}$$

Exercise 0.24. If $T_1, \dots, T_n \in \mathcal{L}(X)$ which commute with each other, then $T_1 \dots T_n$ is invertible if and only if the individual elements are invertible.

We know

$$\begin{aligned} \mu \in \sigma(p(T)) &\iff \mu - p(T) \text{ is not invertible} \\ &\iff T - \lambda I \text{ non invertible for some } i \\ &\iff \lambda \in \sigma(T) \text{ for some } i \\ &\iff \mu = p(\lambda_i) \in p(\sigma(T)) \end{aligned}$$

and so

$$\sigma(p(T)) = p(\sigma(T))$$

□

Definition 0.25 (Spectral radius). Let $X \neq \{0\}$ be a Banach space over \mathbb{C} . The **spectral radius** of $T \in \mathcal{L}(X)$ is

$$\begin{aligned} r(T) &= \sup\{|\lambda| \mid \lambda \in \sigma(T)\} \\ &= \max\{|\lambda| : \lambda \in \sigma(T)\} \end{aligned}$$

Note.

$$r(T) \leq \|T\|$$

since $\sigma(T) \subseteq \{\lambda \in \mathbb{C} \mid |\lambda| \leq \|T\|\}$. Strict inequality can (and often does) occur.

Example 0.26. Let

$$T = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}.$$

Then consider $T : \mathbb{C}^2 \rightarrow \mathbb{C}^2$ where $\|(x, y)\|_2 = \sqrt{|x|^2 + |y|^2}$. Then

$$\begin{aligned} \|T\| &= \sup\{\|Tx\|_2 \mid x \in \mathbb{C}^2\} \\ &= \sqrt{\lambda_{\max}(T^*T)} \end{aligned}$$

where

$$T^* = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}$$

is conjugate transpose. Then $\|T\| = 1$. But $\sigma(T) = \{0\}$, and so $r(T) = 0 < 1 = \|T\|$.

Theorem 0.27 (Gelfand, 1941). *Let $X \neq \{0\}$ be Banach over \mathbb{C} , and let $T \in \mathcal{L}(X)$. Then*

$$r(T) = \lim_{n \rightarrow \infty} \|T^n\|^{1/n}.$$

In particular, the limit exists.

PROOF. By the spectral mapping theorem,

$$\sigma(T^n) = \{\sigma(T)\}^n = \{\lambda^n \mid \lambda \in \sigma(T)\}.$$

So

$$\begin{aligned} r(T) &= r(T^n)^{1/n} \\ &\leq \|T^n\|^{1/n}. \end{aligned}$$

So

$$r(T) \leq \liminf_{n \rightarrow \infty} \|T^n\|^{1/n}$$

Now, we must show that

$$\limsup_{n \rightarrow \infty} \|T^n\|^{1/n} \leq r(T).$$

Let $\varphi \in \mathcal{L}(X)$ and let

$$\begin{aligned} f_\varphi : \rho(T) &\rightarrow \mathbb{C} \\ \lambda &\mapsto \varphi((\lambda I - T)^{-1}) \end{aligned}$$

We saw that f_φ is analytic on $\rho(T)$. We also have

$$f_\varphi(\lambda) = \sum_{n=0}^{\infty} \frac{1}{\lambda^{n+1}} \varphi(T^n) \quad (\star)$$

if $|\lambda| > \|T\|$. By general theory of Laurent series, (\star) actually holds for all $\lambda \in \rho(T)$. In particular, it holds if $|\lambda| > r(T)$.

Thus,

$$\lim_{n \rightarrow \infty} \frac{1}{\lambda^{n+1}} \varphi(T^n) = 0 \quad \boxed{|\lambda| > r(T)}$$

Sp for each $\varphi \in \mathcal{L}(X)'$, and each $|\lambda| > r(T)$, there is $C_{\lambda, \varphi}$ such that

$$\left| \varphi \left(\frac{1}{\lambda^{n+1}} T^n \right) \right| \leq C_{\lambda, \varphi} \quad \forall n \geq 0$$

Then by the principle of uniform boundedness, there exists a constant C_λ such that

$$\left\| \frac{1}{\lambda^{n+1}} T^n \right\| \leq C_\lambda \quad \forall n \geq 0$$

So $\|T^n\|^{1/n} \leq |\lambda|(C_\lambda|\lambda|)^{1/n}$, which gives

$$\limsup_{n \rightarrow \infty} \|T^n\|^{1/n} \leq \lambda$$

for all $|\lambda| > r(T)$. So

$$\limsup_{n \rightarrow \infty} \|T^n\|^{1/n} \leq r(T)$$

We used the following lemma.

Lemma 0.28. *Let X be a normed vector space, $A \subseteq X$ a subset. We say that*

- (1) A is **bounded** if there exists $C > 0$ with $\|x\| \leq C$, for all $x \in A$.
- (2) A is **weakly bounded** if for each $\varphi \in X'$, there exists $C_\varphi > 0$ such that

$$|\varphi(x)| \leq C_\varphi$$

for all $x \in A$.

Then we have

$$A \subseteq X \text{ is bounded} \iff \text{weakly bounded}$$

PROOF. A bounded $\Rightarrow \|x\| \leq C$ for all $x \in A \Rightarrow |\varphi(x)| \leq \|\varphi\|\|x\| \leq \|\varphi\|C$. So A is weakly bounded.

Now, suppose A is weakly bounded. For each $x \in X$, let $\hat{x} \in X''$ with

$$\hat{x}(\varphi) = \varphi(x).$$

So $|\hat{x}(\varphi)| \leq C_\varphi$ for all $x \in A$. By the principle of uniform boundedness,

$$\|\hat{x}\| \leq C$$

for all $x \in A$, and since $\|\hat{x}\| = \|x\|$. Thus A is bounded. □

□

Compact Operators

We now turn to compact operators. In general, calculating $\sigma(T)$ is difficult, but for compact operators on a complex Banach space, we have a fairly explicit theory.

Theorem 0.29. *Let X be a complex Banach space, with $\dim(X) = \infty$. Let $T : X \rightarrow X$ be a compact operator. Then*

- (1) $0 \in \sigma(T)$.
- (2) $\sigma(T) \setminus \{0\} = \sigma_p(T) \setminus \{0\}$, that is, each $\lambda \in \sigma(T) \setminus \{0\}$ is an eigenvalue of T (0 may or may not be an eigenvalue.)
- (3) We are in exactly one of the cases:
 - $\sigma(T) = \{0\}$.
 - $\sigma(T) \setminus \{0\}$ is finite (nonempty).
 - $\sigma(T) \setminus \{0\}$ is a sequence of points converging to 0.
- (4) Each $\lambda \in \sigma(T) \setminus \{0\}$ is isolated, and the eigenspace $\text{KER}(\lambda I - T)$ is finite dimensional.

where $\sigma_p(T)$ is the **point spectrum of T** , where

$$\begin{aligned} \sigma_p(T) &= \{\lambda \in \mathbb{K} \mid \lambda I - T \text{ is not injective}\} \\ &= \{\lambda \in \mathbb{K} \mid \text{there exists nonzero vector } x \text{ with } (\lambda I - T)x = 0\} \\ &= \{\text{eigenvalues of } T\} \end{aligned}$$

PROOF. We shall prove these results next week. □

Definition 0.30. Let X, Y be normed vector spaces. An operator $T : X \rightarrow Y$ is **compact** if T is linear, and if $B \subseteq X$ is bounded then $T(B)$ is relatively compact (a set is relatively compact if its closure is compact.) Symbolically,

$$B \subseteq X \text{ bounded} \Rightarrow \overline{T(B)} \text{ compact}$$

Lemma 0.31. *If T is compact, then T is continuous.*

PROOF. The closed ball $B = \{x \in X \mid \|x\| \leq 1\}$ is bounded, and so if T is a compact operator, then $\overline{T(B)}$ is compact, and hence bounded. Hence $\|Tx\| \leq M$ for all $\|x\| \leq 1$, so T is continuous, with $\|T\| \leq M$. □

We now recall definitions of compactness

Theorem 0.32 (Characterisations of compactness). *Let X be a metric space. The following are equivalent.*

- (1) X is **compact** (every open cover has a finite subcover).
- (2) X is **sequentially compact** (every sequence in X has a convergent subsequence)

Lemma 0.33. *Let X be a compact set. Let $Y \subseteq X$. If $Y \subseteq X$ is closed, then Y is compact.*

Lemma 0.34. *Let V be a finite dimensional vector space. If $X \subseteq V$ is closed and bounded, then X is compact.*

Theorem 0.35 (Characterisations of compact operators). *Let X, Y be normed vector spaces over \mathbb{K} . Let $T \in \mathcal{L}(X, Y)$. Then the following are equivalent.*

- (a) T is compact.
- (b) $\overline{T(B)}$ is compact, where $B = \{x \in X \mid \|x\| \leq 1\}$.
- (c) If $(x_n)_{n \geq 1}$ is bounded in X , then $(Tx_n)_{n \geq 1}$ has a convergent subsequence (**sequentially compact**).

PROOF. (a) \Rightarrow (b) by definition.

(b) \Rightarrow (a). Suppose (b) holds. Let $B_1 \subseteq X$ be bounded. Then $B_1 \subseteq \alpha B$ for some $\alpha > 0$. So

$$\overline{T(B_1)} \subseteq \overline{T(\alpha B)} = \alpha \overline{T(B)}$$

which is a closed subset of a compact set, and hence compact.

(a) \Rightarrow (c). Suppose T is compact. Let $(x_n)_{n \geq 1}$ be bounded sequence in X . Then $T(B) = \{Tx_n \mid n \geq 1\}$ is relatively compact. So $\overline{T(B)}$ is compact, and hence is sequentially compact, and so has a convergence subsequence.

(c) \Rightarrow (a). Let $B \subseteq X$ be bounded. Let $(y_n)_{n \geq 1}$ be a sequence in $T(B)$. Then there is $x_n \in B$ with $Tx_n = y_n$. So $(x_n)_{n \geq 1}$ is a bounded sequence. By assumption $(Tx_n)_{n \geq 1}$ has a convergent subsequence. So $\overline{T(B)}$ is sequentially compact, and hence compact. \square

Corollary. *The set $\{\text{compact operators } T : X \rightarrow Y\}$ is a vector space. That is, if T_1, T_2 are compact, then $T_1 + T_2$ and αT_1 are compact.*

PROOF. Exercise. Use (c) from the characterisation of compact operators. \square

Corollary.

$$\mathcal{K}(X, Y) \subseteq \mathcal{L}(X, Y) \subseteq \text{Hom}(X, Y)$$

where $\mathcal{K}(X, Y)$ is the set of compact operators $T : X \rightarrow Y$.

Example 0.36 (Finite rank operators). Let X, Y be normed vector spaces, and let $T \in \mathcal{L}(X, Y)$. If $\dim(\text{Im } T) < \infty$, then T is said to have **finite rank**. Then if T has finite rank, then T is compact.

PROOF. Let (x_n) be a bounded sequence in X . Then $\|Tx_n\| \leq \|T\|\|x_n\|$ so (Tx_n) is a bounded sequence in $\text{Im } T$. But $\text{Im } T$ is finite dimensional, and so $\overline{\{Tx_n \mid n \geq 1\}}$ is compact (closed and bounded), and so $(Tx_n)_{n \geq 1}$ has a convergent subsequence. By (c) in Theorem 0.35, T is compact. \square

Lemma 0.37. *Let X, Y be normed vector spaces. If $T \in \mathcal{L}(X, Y)$ has finite rank, then there exists $y_1, \dots, y_n \in \text{Im } T$ and $\varphi_1, \dots, \varphi_n \in X'$ with $Tx = \sum_{j=1}^n \varphi_j(x)y_j$ for all $x \in X$, with $n = \dim(\text{Im } T)$.*

PROOF. Choose a basis y_1, \dots, y_n of $\text{Im } T$. For each $j = 1, \dots, n$, define $\alpha_j \in (\text{Im } T)'$ by

$$\alpha_j(a_1y_1 + \dots + a_ny_n) = a_j$$

i.e. coordinate projection. By Hahn-Banach, we can extend α_j to a continuous linear functional $\tilde{\alpha}_j \in Y'$. Let $\varphi_j = \tilde{\alpha}_j \circ T : X \rightarrow \mathbb{K}$. So $\varphi_j \in X'$. Since

$$y = \sum_{j=1}^n \tilde{\alpha}_j(y)y_j \quad \forall y \in \text{Im } T$$

we have

$$\begin{aligned} Tx &= \sum_{j=1}^n \tilde{\alpha}_j(Tx)y_j \\ &= \sum_{j=1}^n (\alpha_j \circ T)(x)y_j \\ &= \sum_{j=1}^n \varphi_j(x)y_j \quad \forall x \in X. \end{aligned}$$

\square

Recall that the closed unit ball in X is compact if and only if $\dim(X) < \infty$. Then it follows that the identity map $I : X \rightarrow X$ is compact if and only if $\dim(X) < \infty$. Hence,

$$\mathcal{K}(X) \subsetneq \mathcal{L}(X) \subsetneq \text{Hom}(X, X)$$

when $\dim(X) = \infty$.

Consider a sequence of compact operators T_n . If T_n is compact and $T_n \rightarrow T$, then T is compact.

Lemma 0.38 (Riesz's Lemma). *Let X be a normed vector space. Let $Y \subsetneq X$ be a proper **closed** subspace. Let $\theta \in (0, 1)$ be given. Then there exists x with $\|x\| = 1$ such that $\|x - y\| \geq \theta$ for all $y \in Y$.*

PROOF. Pick any $z \in X \setminus Y$. Let $\alpha = \inf_{y \in Y} \|z - y\| > 0$ since Y is closed. Then by the definition of the infimum, there is $y_0 \in Y$ with $\alpha \leq \|z - y_0\| \leq \frac{\alpha}{\theta}$. Now let $x = \frac{z - y_0}{\|z - y_0\|}$. Then $\|x\| = 1$.

Now,

$$\begin{aligned}\|x - y\| &= \left\| \frac{z - y_0}{\|z - y_0\|} - y \right\| \\ &= \frac{1}{\|z - y_0\|} \|z - y_0 - \|z - y_0\|y\| \\ &\geq \frac{\theta}{\alpha} = \theta\end{aligned}$$

□

Corollary. *Let X be a normed vector space. The closed unit ball $\overline{B}(0,1)$ is compact if and only if $\dim(X) < \infty$.*

PROOF. If $\dim(X) < \infty$ then $\overline{B}(0,1)$ is compact (since closed and bounded if and only if compact in finite dimensions). Now suppose $\dim(X) = \infty$. Build a sequence (x_n) with $\|x_n\| = 1$ with no convergent subsequence. Choose finite dimensional subspaces

$$\{0\} = X_0 \subsetneq X_1 \subsetneq X_2 \subsetneq \dots$$

These are all closed (finite dimensional spaces are complete, and hence closed). Use the lemma to choose $x_k \in X_k$ with $\|x_k\| = 1$, $\|x_k - x\| \geq \frac{1}{2}$ for all $x \in X_{k-1}$. So $\|x_k - x\| \geq \frac{1}{2}$ for all $x \in X_j (j \leq k-1)$. So $\|x_n - x_m\| \geq \frac{1}{2}$ for all $m, n \geq 1$. So (x_n) has no convergent subsequence, and so $\overline{B}(0,1)$ is not compact. □

Corollary. *$I : X \rightarrow X$ is compact if and only if $\dim(X) < \infty$.*

PROOF. Recall T is compact if and only if $T(\overline{B}(0,1))$ is relatively compact. □

One way to show that an operator is compact is to apply the following.

Proposition 0.39. *Let X be a normed vector space, and let Y be Banach. Suppose that $T_n \in \mathcal{K}(X, Y)$ for each $n \geq 1$. If $T_n \rightarrow T$ (in operator norm, $\|T_n - T\| \rightarrow 0$) then T is compact.*

PROOF. Let (x_n) be a bounded sequence in X . We now construct a subsequence (x'_n) for which (Tx'_n) converges.

- Since T_1 is compact, (x_n) has a subsequence $x_n^{(1)}$ such that $(T_1 x_n^{(1)})$ converges.
- Since T_2 is compact and $x_n^{(1)}$ is bounded, there is a subsequence $x_n^{(2)}$ such that $T_2 x_n^{(2)}$ converges.
- Continuing, we can form a subsequence $x_n^{(k)}$ such that $T_k x_n^{(k)}$ converges.

Let $x'_n = x_n^{(n)}$. Then (x'_n) is a subsequence of $(x_n^{(1)})$, and $(x'_n)_{n \geq 2}$ is a subsequence of $(x_n^{(2)})$, etc. So for each fixed $k \geq 1$, $(T_k x'_n)$ converges.

We now show Tx'_n is Cauchy, and hence converges. We have

$$\|Tx'_m - Tx'_n\| \leq \|Tx'_m - T_k x'_m\| + \|T_k x'_m - T_k x'_n\| + \|T_k x'_n - Tx'_n\|$$

where k is to be chosen. Suppose $\|x_n\| \leq M$ for all $n \geq 1$. Then

$$\|Tx'_m - Tx'_n\| \leq 2M\|T - T_k\| + \|T_kx'_m - T_kx'_n\|$$

Let $\epsilon > 0$ be given. Since $\|T - T_k\| \rightarrow 0$ as $k \rightarrow \infty$, fix a k for which $\|T - T_k\| \leq \frac{\epsilon}{3M}$. For this fixed k , we know $(T_kx'_n)$ converges, and so is Cauchy. So there exists $N < \infty$ such that $\|T_kx'_m - T_kx'_n\| < \epsilon$ for all $m, n < N$. Hence $\|Tx'_m - Tx'_n\| \leq \frac{2M}{\epsilon}3M + \frac{\epsilon}{3} = \epsilon$ for all $m, n > N$, so is Cauchy, and so converges. \square

Example 0.40. Let $K(x, y) \in L^2(\mathbb{R}^2)$. Define $T : L^2(\mathbb{R}) \rightarrow L^2(\mathbb{R})$ by

$$Tf(x) = \int_{\mathbb{R}} K(x, y)f(y) dy$$

(Hilbert-Schmidt Integral operator)

Proposition 0.41. T is compact.

PROOF. Note that $\|Tf\|_2 \leq \|K\|_2\|f\|_2$ for all $f \in L^2(\mathbb{R})$, where $\|K\|_2 = (\iint_{\mathbb{R}^2} |K(x, y)|^2 dx dy)^{1/2}$. So T is continuous, with $\|T\| \leq \|K\|_2$. We now exhibit T as a limit of finite rank (hence compact) operators, with $T_n : L^2(\mathbb{R}) \rightarrow L^2(\mathbb{R})$. One can see that there is a sequence $K_n \in L^2(\mathbb{R}^2)$ of the form

$$K_n(x, y) = \sum_{k=1}^{N_n} \alpha_k^{(n)}(x)\beta_k^{(n)}(y)$$

with $K_n \rightarrow K$ in $L^2(\mathbb{R}^2)$. Then $\|T_n - T\| \leq \|K_n - K\|_2 \rightarrow 0$, and so $T_n \rightarrow T$. Hence

$$\begin{aligned} T_n f(x) &= \sum_{k=1}^{N_n} \int_{\mathbb{R}} \alpha_k^{(n)}(x)\beta_k^{(n)}(y)f(y) dy \\ &= \sum_{k=1}^{N_n} \langle f, \overline{\beta_k^{(n)}} \rangle \alpha_k^{(n)}(x) \end{aligned}$$

and so $T_n f = \sum_{k=1}^{N_n} \langle f, \overline{\beta_k^{(n)}} \rangle \alpha_k^{(n)}$ from which we use that T_n has finite rank. \square

Theorem 0.42. Let X be a complex Banach space, with $\dim(X) = \infty$. Let $T : X \rightarrow X$ be a compact operator. Then

- (1) $0 \in \sigma(T)$.
- (2) $\sigma(T) \setminus \{0\} = \sigma_p(T) \setminus \{0\}$, that is, each $\lambda \in \sigma(T) \setminus \{0\}$ is an eigenvalue of T (0 may or may not be an eigenvalue.)
- (3) We are in exactly one of the cases:
 - $\sigma(T) = \{0\}$.
 - $\sigma(T) \setminus \{0\}$ is finite (nonempty).
 - $\sigma(T) \setminus \{0\}$ is a sequence of points converging to 0.
- (4) Each $\lambda \in \sigma(T) \setminus \{0\}$ is isolated, and the eigenspace $\text{KER}(\lambda I - T)$ is finite dimensional.

where $\sigma_p(T)$ is the **point spectrum of T** , where

$$\begin{aligned}\sigma_p(T) &= \{\lambda \in \mathbb{K} \mid \lambda I - T \text{ is not injective}\} \\ &= \{\lambda \in \mathbb{K} \mid \text{there exists nonzero vector } x \text{ with } (\lambda I - T)x = 0\} \\ &= \{\text{eigenvalues of } T\}\end{aligned}$$

Compact operators are very well behaved with respect to composition.

Proposition 0.43. *Let X, Y, Z be normed vector spaces.*

(a) *If $T \in \mathcal{K}(X, Y)$ and $S \in \mathcal{L}(Y, Z)$, then $ST \in \mathcal{K}(X, Z)$.*

(b) *If $S \in \mathcal{L}(X, Y)$ and $T \in \mathcal{K}(Y, Z)$, then $TS \in \mathcal{K}(X, Z)$.*

PROOF. (a) Let (x_n) be a bounded sequence in X . Since T is compact, Tx_n has a convergent subsequence, say $Tx_{n_k} \rightarrow y \in Y$. Then (STx_n) has a convergent subsequence, namely $STx_{n_k} = S(Tx_{n_k}) \rightarrow Sy$ by continuity of S . So ST is compact.

(b) Let $B \subseteq X$ be bounded. Then $S(B)$ is bounded in Y , as S is continuous. So $TS(B) = T(S(B))$ is relatively compact since T is compact. Hence TS is compact. \square

Corollary (Part (1) of theorem). *If X is infinite dimensional Banach space, then $0 \in \sigma(T)$.*

PROOF. If $0 \notin \sigma(T)$ then T is invertible. By bounded inverse theorem T^{-1} is continuous, and then $I = TT^{-1}$ is compact, which is a contradiction. \square

Theorem 0.44 (Part (3) of theorem). *Let X be a normed vector space. Let $T \in \mathcal{K}(X)$. Then T has at most countably many eigenvalues. If T has infinitely many eigenvalues, then they can be arranged in a sequence converging to zero.*

PROOF. We show that for each $N > 0$, we have

$$\#\{\lambda \in \sigma_p(T) \mid |\lambda| \geq N\} < \infty \quad (\star)$$

Suppose that there is $N > 0$ such that (\star) fails. So $\lambda_1, \lambda_2, \dots$ are distinct eigenvalues with $|\lambda_n| \geq N$ for $n = 1, 2, \dots$. Let $x_n \neq 0$ be an eigenvector. $Tx_n = \lambda_n x_n$, $n = 1, 2, \dots$. Let $X_n = \text{span}\{x_1, \dots, x_n\}$. Since $\{x_n \mid |x_n| \geq 1\}$ are linearly independent, we have

$$X_1 \subsetneq X_2 \subsetneq \dots$$

and each X_n is closed (finite dimensional).

By Reisz's Lemma from previous lecture, choose $y_n \in X_n$ such that $\|y_n\| = 1$ and $\|y_n - x\| \geq \frac{1}{2}$ for all $x \in X_{n-1}$. So (y_n) is bounded in X . We show that Ty_n has no convergence subsequence, contradicting compactness of T .

Let $m > n$. Then

$$\begin{aligned} \|Ty_m - Ty_n\| &= \|\lambda_m y_m - (\lambda_m y_m - Ty_m + Ty_n)\| \\ &= |\lambda_m| \|y_m - (\text{something in } X_{m-1})\| \\ &\geq \frac{1}{2} |\lambda_m| \geq \frac{1}{2} N \end{aligned}$$

as required.

Note that $y_m = a_1 x_1 + \cdots + a_m x_m$. Then

$$\begin{aligned} \lambda_m y_m - Ty_m &= \lambda_m a_1 x_1 + \cdots + \lambda_m a_m x_m - (a_1 \lambda_1 x_1 + \cdots + a_m \lambda_m x_m) \\ &= a_1 (\lambda_m - \lambda_1) x_1 + \cdots + a_{m-1} (\lambda_m - \lambda_{m-1}) x_{m-1} \in X_{m-1} \end{aligned}$$

and $Ty_n \in X_{m-1}$ since $n < m$. □

Definition 0.45 (Projection operator). Let X be a vector space. A linear operator $P : X \rightarrow X$ is called a projection if $P^2 = P$.

Proposition 0.46. *If $P : X \rightarrow X$ is a projection then $I - P$ is a projection, and*

$$\text{IM } I - P = \text{KER } P, \quad \text{KER } I - P = \text{IM } P$$

PROOF. If $P^2 = P$ then $(I - P)^2 = I - 2P + P^2 = I - P$ and so $I - P$ is a projection. Let $x \in \text{IM } I - P$. Then $x = (I - P)y$ for some $y \in X$. So $Px = P(I - P)y = (P - P^2)y = 0$. So $x \in \text{KER } P$ and $\text{IM } I - P \subseteq \text{KER } P$. If $x \in \text{KER } P$ then $Px = 0$. So $(I - P)x = x$, and $x \in \text{IM } (I - P)$. □

Definition 0.47 (Direct sum). Let X be a vector space, and let X_1, X_2 be subspaces. Then $X = X_1 \oplus X_2$ (direct sum) if

$$X = X_1 + X_2$$

and $X_1 \cap X_2 = \{0\}$. Equivalently, $X = X_1 \oplus X_2$ if and only if each $x \in X$ can be written in exactly one way as $x = x_1 + x_2$ with $x_1 \in X_1, x_2 \in X_2$.

Theorem 0.48 (Equivalence of direct sums and projections). *Let X be a vector space.*

(a) *If $P : X \rightarrow X$ is a projection, then*

$$X = (\text{IM } P) \oplus (\text{KER } P)$$

(b) *If $X = X_1 \oplus X_2$, there exists a unique projection with*

$$\text{IM } P = X_1, \quad \text{KER } P = X_2.$$

Specifically, $Px = x_1$ if $x = x_1 + x_2$.

PROOF. (a) Let $P : X \rightarrow X$ be a projection. Then we show $X = (\text{IM } P) \oplus (\text{IM } I - P)$, $x = Px + (I - P)x$. This shows that $X = \text{IM } P + \text{IM } I - P$. If $x \in \text{IM } P \cap \text{KER } P$ then $x = Py$ and $Px = 0$. Hence, $Px = P^2y = P^y = 0$ and so $x = 0$.

(b) Exercise. □

Proposition 0.49. *Let X be Banach. Let $X = X_1 \oplus X_2$. Let $P : X \rightarrow X$ be the corresponding projection operator. Then*

$$P \in \mathcal{L}(X) \iff X_1, X_2 \text{ closed}$$

PROOF. (\Rightarrow). Suppose P is continuous. Then $X_1 = \text{IM } P = \text{KER } I - P$ and $X_2 = \text{KER } P$ are both closed. For example, if $x_n \in \text{KER } P$ and $x_n \rightarrow x$, then $0 = Px_n \rightarrow Px$ and so $x \in \text{KER } P$.

(\Leftarrow). Suppose that X_1, X_2 are closed. Since $X = X_1 \oplus X_2$, we can define a new norm $\|\cdot\|'$ by $\|x\|' = \|x_1\| + \|x_2\|$ where $x = x_1 + x_2$.

Exercise 0.50.

(a) Show that $\|\cdot\|'$ is a norm.

(b) Show that $(X, \|\cdot\|')$ is Banach. This relies on the fact that $(X, \|\cdot\|)$ is Banach and X_1, X_2 are closed.

Note that $\|x\| = \|x_1 + x_2\| \leq \|x_1\| + \|x_2\| = \|x\|'$, and so by a corollary to the open mapping theorem, there is a $c > 0$ with $\|x\|' \leq c\|x\|$ for all $x \in X$, and so

$$\|Px\| = \|x_1\| \leq \|x_1\| + \|x_2\| = \|x\|' \leq c\|x\|$$

and hence P is continuous. □

Corollary. *Let X be Banach, and let M be a finite dimensional subspace. Then there exists a closed N with*

$$X = M \oplus N.$$

PROOF. Let v_1, \dots, v_n be a basis of M . Define, for each $j = 1, \dots, n$, $\varphi_j \in M'$ by $\varphi_j(a_1v_1 + \dots + a_nv_n) = a_j$. Then using Hahn-Banach to extend $\tilde{\varphi}_j \in X'$. Let $P : X \rightarrow X$ be defined by

$$Px = \sum_{j=1}^n \tilde{\varphi}_j(x)v_j.$$

Then we need only check that P is linear and continuous, $\text{IM } P = M$, and $P^2 = P$. Now take $N = \text{KER } P$ and then $X = M \oplus N$. □

.

We are now ready to prove the following theorem.

Theorem 0.51. *Let X be Banach, and let $T \in \mathcal{K}(X)$, and let $\lambda \in \mathbb{K} \setminus \{0\}$. For all $k \in \mathbb{N}$, we have*

- (a) $\underbrace{\text{KER } (\lambda I - T)^k}_{\text{generalised eigenspace}}$ is finite dimensional.
- (b) $\text{IM } (\lambda I - T)^k$ is closed.

PROOF. Reductions. Since $\text{KER } (\lambda I - T)^k = \text{KER } (I - \lambda^{-1}T)^k$, and similarly for the image, by replacing $T \in \mathcal{K}(X)$ by $\lambda T \in \mathcal{K}(X)$, we can assume that $\lambda = 1$.

Also, we have

$$\begin{aligned} (I - T)^k &= \sum_{n=0}^k \binom{k}{n} (-1)^n T^n \\ &= I - T \underbrace{\sum_{n=1}^k \binom{k}{n} (-1)^{n-1} T^{n-1}}_{\text{continuous}} \\ &= I - \tilde{T}. \end{aligned}$$

where \tilde{T} is the composition of compact and continuous operators, and so is compact. So we can take $\lambda = 1, k = 1$.

- (a) The closed unit ball in $\text{KER } I - T$ is

$$\begin{aligned} \{x \in \text{KER } I - T \mid \|x\| \leq 1\} &= \{Tx \mid x \in \text{KER } I - T, \|x\| \leq 1\} \\ &\subseteq \overline{T(\overline{B(0, 1)})} \end{aligned}$$

which is compact as T is compact. Hence, the closed unit ball in $\text{KER } I - T$ is compact, and thus $\text{KER } I - T$ is finite dimensional.

- (b) Let $S = I - T$. We then need to show that $\text{IM } S$ is closed. Since $\text{KER } S$ is finite dimensional from above, there is a **closed** subspace N with

$$X = (\text{KER } S) \oplus N$$

Note that $\text{IM } S = S(X) = S(N)$, and that $S|_N : N \rightarrow X$ is injective.

Suppose that $S(N)$ is not closed. So there is a sequence (x_n) in N such that $Sx_n \rightarrow y \in X \setminus S(N)$. Then there are two cases

Case 1 ($\|x_n\| \rightarrow \infty$). Let $y_n = \frac{1}{\|x_n\|} x_n$. Then $Sy_n = \frac{1}{\|x_n\|} Sx_n \rightarrow 0$. But $(y_n)_{n \geq 1}$ is bounded in X , and so there exists a subsequence y_{n_k} such that $Ty_{n_k} \rightarrow z$ (as T is compact). Hence $y_{n_k} = Sy_{n_k} + Ty_{n_k} \rightarrow 0 + z$. Thus $z \in N$ (as $y_{n_k} \in N$, and N is closed), and $\|z\| = 1$.

So $Sy_{n_k} \rightarrow 0$, but $Sy_{n_k} \rightarrow Sz$ with $z \in N \setminus \{0\}$, by the continuity of S . This contradicts the injectivity of $S|_N$.

Case 2 ($\|x_n\|$ does not tend to infinity). So (x_n) has a bounded subsequence (x_{n_k}) . Since T is compact, (x_{n_k}) has a subsequence such that $(Tx_{n_{k_l}})$ converges, to z_1 say. By replacing x_n

by this subsequence we can assume that $Sx_n \rightarrow y$, and that $Tx_n \rightarrow z$. A before, we can write

$$x_n = Sx_n + Tx_n \rightarrow y + z.$$

So x_n converges to $x \in N$. So $Sx_n \rightarrow Sx \in S(N)$ by continuity, but we assume that $Sx_n \rightarrow y \in X \setminus S(N)$, which achieves our contradiction.

□

Let $T : \mathbb{C}^n \rightarrow \mathbb{C}^n$ be a linear operator. Then in the simplest case, T has n distinct eigenvalues, and the corresponding eigenvectors are linearly independent, forming a basis for \mathbb{C}^n .

Hence, $\mathbb{C}^n = \mathbb{C}x_1 \oplus \cdots \oplus \mathbb{C}x_n$ and the matrix of T relative to this basis is simply diagonal with $\lambda_1, \dots, \lambda_n$.

This is not always possible, because there is not always a basis of eigenvectors. Instead look at the generalised eigenspace,

$$\{x \in \mathbb{C}^n \mid (\lambda I - T)^k x = 0 \text{ for some } k \geq 1.\}$$

But $\{0\} \subseteq \text{KER}(\lambda I - T)^1 \subseteq \text{KER}(\lambda I - T)^2 \subseteq \dots$ and since $\dim(\mathbb{C}^n) < \infty$ this must stabilise. Let $r \geq 1$ be the first time that $\text{KER}(\lambda I - T)^r = \text{KER}(\lambda I - T)^{r+1}$. Then the generalised λ -eigenspace is just $\text{KER}(\lambda I - T)^r$. There is a basis of \mathbb{C}^n consisting of generalised eigenvectors, and the matrix of T relative to this basis is in block form.

Definition 0.52 (Complete reduction). Let $T : X \rightarrow X$ be linear. If $X = X_1 \oplus X_2$ we can write

$$Tx = \begin{pmatrix} T_{11} & T_{12} \\ T_{21} & T_{22} \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix}$$

where we identify $x_1 + x_2 \iff (x_1, x_2)$. Here,

$$T_{11} : X_1 \rightarrow X_1$$

$$T_{12} : X_2 \rightarrow X_1$$

$$T_{21} : X_2 \rightarrow X_2$$

$$T_{22} : X_2 \rightarrow X_2$$

we say that $X = X_1 \oplus X_2$ **completely reduces** T (well adapted to T) if

$$Tx = \begin{pmatrix} T_1 & 0 \\ 0 & T_2 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix}$$

We write $T = T_1 \oplus T_2$.

Exercise 0.53. If $X = X_1 \oplus X_2$ completely reduces $T = T_1 \oplus T_2$, then

(a) $\text{KER } T = \text{KER } T_1 \oplus \text{KER } T_2$

(b) $\text{IM } T = \text{IM } T_1 \oplus \text{IM } T_2$

- (c) T is injective if and only if T_1, T_2 are injective
- (d) T is surjective if and only if T_1, T_2 are surjective
- (e) If T is bijective, then $X = X_1 \oplus X_2$ completely reduces $T^{-1} = T_1^{-1} \oplus T_2^{-1}$.

Corollary. *Let $X = X_1 \oplus X_2$ be Banach, with X_1, X_2 closed subspaces. If $X = X_1 \oplus X_2$ completely reduces $T = T_1 \oplus T_2 \in \mathcal{L}(X)$, then*

- (a) $T_1 \in \mathcal{L}(X_1), T_2 \in \mathcal{L}(X_2)$
- (b) $\sigma(T) = \sigma(T_1) \cup \sigma(T_2)$
- (c) $\sigma_p(T) = \sigma_p(T_1) \cup \sigma_p(T_2)$

PROOF. Exercise. □

Consider the following chains

$$\begin{aligned} \{0\} &\subseteq \text{KER } S^1 \subseteq \text{KER } S^2 \subseteq \dots \\ X &\supseteq \text{IM } S^1 \supseteq \text{IM } S^2 \supseteq \dots \end{aligned}$$

where X is a vector space and $S \in \text{Hom}(X, X)$. It is easy to see that if $\text{KER } S^r = \text{KER } S^{r+1}$ then $\text{KER } S^r = \text{KER } S^{r+k}$. Similarly for images (p. 109 in Daners.)

There is no reason that these should stabilise in general.

Theorem 0.54. *Let X be Banach, $T \in \mathcal{K}(X), \lambda \neq 0$. Then both chains (with $S = \lambda I - T$) stabilise.*

PROOF. Without loss of generality, assume $\lambda = 1$, so we can write $S = I - T$. Suppose that the kernel chain does not stabilise. Since we assume

$$\text{KER } S^1 \subsetneq \text{KER } S^2 \subsetneq \text{KER } S^3 \subsetneq \dots$$

We know that these are closed (being finite dimensional) subspaces. So Reisz's Lemma gives $x_n \in \text{KER } S^n$ with $\|x_n\| = 1, \|x_n - x\| \geq \frac{1}{2}$ for all $x \in \text{KER } S^{n+1}$. This is a bounded sequence. We claim that Tx_n has no convergent subsequence.

Let $m > n$. Then

$$\begin{aligned} \|Tx_m - Tx_n\| &= \|(I - T)x_n - (I - T)x_m + x_m - x_n\| \\ &= \|Sx_n - Sx_m - x_m - x_n\| \\ &= \|x_m - \underbrace{(Sx_m - Sx_n + x_n)}_{\text{in KER } S^{m-1}}\| \\ &\geq \frac{1}{2} \end{aligned}$$

The image argument is similar - using the fact that the images are closed - proved in the previous lecture. □

Theorem 0.55. *Let X be a vector space, $S \in \text{Hom}(X, X)$. Suppose that*

$$\alpha(S) = \inf\{r \geq 1 \mid \text{KER } S^r = \text{KER } S^{r+1}\}$$

$$\delta(S) = \inf\{r \geq 1 \mid \text{IM } S^r = \text{IM } S^{r+1}\},$$

*the **ascent** and **descent** of S respectively, are both finite.*

Then

(a) $\alpha(S) = \delta(S) = r$, say

(b) $X = \text{KER } S^r \oplus \text{IM } S^r$

(c) *The direct sum in (b) completely reduces S .*

PROOF. Daner's notes, p. 109. □

Corollary. *Let X be Banach, $T \in \mathcal{K}(X)$, $\lambda \neq 0$. Let $r = \alpha(\lambda I - T) = \delta(\lambda I - T)$. Then $X = \text{KER } (\lambda I - T)^r \oplus \text{IM } (\lambda I - T)^r$ and this completely reduces $\mu I - T, \mu \in \mathbb{K}$.*

Corollary. *If X is Banach, $T \in \mathcal{K}(X), \lambda \neq 0$ then $\lambda I - T$ is injective if and only if $\lambda I - T$ is surjective.*

PROOF.

$$\begin{aligned} & \lambda I - T \text{ injective} \\ \Rightarrow & 0 \in \text{KER } (\lambda I - T)^1 = \text{KER } (\lambda I - T)^2 \\ \Rightarrow & \alpha(\lambda I - T) = 1 \\ \Rightarrow & \delta(\lambda I - T) = 1 \\ \Rightarrow & X = \underbrace{\text{KER } (\lambda I - T)}_{=\{0\}} \oplus \text{IM } (\lambda I - T) \\ \Rightarrow & X = \text{IM } (\lambda I - T) \\ \Rightarrow & X \text{ is surjective} \end{aligned}$$

The other direction is similar. □

Corollary. *Let X be Banach, $T \in \mathcal{K}(X)$. Thus each $\lambda \in \sigma(T) \setminus \{0\}$ is an eigenvalue.*

PROOF. Immediate from the previous corollary. □

The Hilbert Space Decomposition

Recall the following.

Corollary. *Let X be Banach, $T \in \mathcal{K}(X)$, $\lambda \neq 0$. Let $r = \alpha(\lambda I - T) = \delta(\lambda I - T)$. Then $X = \text{KER } (\lambda I - T)^r \oplus \text{IM } (\lambda I - T)^r$ and this completely reduces $\mu I - T, \mu \in \mathbb{K}$.*

Also note that $\text{IM KER } (\lambda I - T)^r$ is closed, and $\text{KER } (\lambda I - T)^r$ is finite dimensional.

Exercise 0.56. Let $\lambda_1, \dots, \lambda_n \in \sigma(T) \setminus \{0\}$. Let $N_j = \text{KER } (\lambda_j I - T)_j^r$ be the generalised λ_j -eigenspace. Show that there exists closed subspaces M with

$$X = N_1 \oplus N_2 \oplus \dots \oplus M$$

with $T = T_1 \oplus T_2 \oplus \dots \oplus T_M$, and so spectral theory tells us how to *diagonalise* T .

In Hilbert spaces we can say even more. Recall that the adjoint of $T \in \mathcal{L}(\mathcal{H})$ is defined by

$$\langle Tx, y \rangle = \langle x, T^*y \rangle \quad \forall x, y \in \mathcal{H}$$

Then $T^* \in \mathcal{L}(\mathcal{H})$.

Definition 0.57 (Self-adjoint). $T \in \mathcal{L}(\mathcal{H})$ is

- (a) **Hermitian (self-adjoint)** if $T^* = T$.
- (b) **Unitary** if $T^*T = TT^* = I$.
- (c) **Normal** if $T^*T = TT^*$.

Remark. For matrices, we have

- (a) Hermitian if and only if $\overline{A^T} = A$.
- (b) Unitary if and only if the columns of A are orthonormal.
- (c) Hermitian and unitary operators are normal.

Proposition 0.58. *Let \mathcal{H} be Hilbert over \mathbb{C} . IF $T \in \mathcal{L}(\mathcal{H})$ is normal, then $r(T) = \|T\|$.*

PROOF. For Hermitian operators it is easy. We have

$$\|T\|^2 = \|T^*T\| = \|T^2\|.$$

By induction, we then have $\|T\|^{2^n} = \|T^{2^n}\|$. So

$$\begin{aligned} r(T) &= \lim_{n \rightarrow \infty} \|T^n\|^{1/n} \\ &= \lim_{n \rightarrow \infty} \|T^{2^n}\|^{1/2^n} \\ &= \|T\|. \end{aligned}$$

For normal operators, we have

$$\begin{aligned} \|T^2\|^2 &= \|(T^2)^*T^2\| \\ &= \|T^*(T^*T)T\| \\ &= \|T^*TT^*T\| \quad \text{normal} \\ &= \|(T^*T)^*(T^*T)\| \\ &= \|T^*T\|^2 \\ &= \|T^4\| \end{aligned}$$

and then we have $\|T^2\| = \|T\|^2$ and the proof follows by induction. \square

Corollary. *Let \mathcal{H} be a Hilbert space over \mathbb{C} .*

(a) *If $T \in \mathcal{L}(\mathcal{H})$ is unitary, then*

$$\sigma(T) \subseteq \mathbb{T} = \{\lambda \in \mathbb{C} \mid |\lambda| = 1\}$$

(b) *If $T \in \mathcal{L}(\mathcal{H})$ is Hermitian, then*

$$\sigma(T) \subseteq \mathbb{R}.$$

PROOF.

(a) On practice sheet. Use the fact that $\sigma(T^*) = \overline{\sigma(T)}$.

(b) Let $\lambda = a + ib \in \sigma(T)$. So $\lambda I - T$ is not invertible. Hence, $(\lambda + it)I - (T + itI)$ is not invertible for all $t \in \mathbb{R}$. Then

$$\begin{aligned} \|\lambda + it\|^2 &\leq r(T + itI)^2 \\ &\leq \|T + itI\|^2 \\ &= \|(T + itI)^*(T + itI)\| \\ &= \|(T - itI)(T + itI)\| \\ &= \|T^2 + t^2I\| \\ &\leq \|T^2 + t^2\| \end{aligned}$$

However, the left hand side is equal to

$$a^2 + b^2 + 2bt + t^2,$$

and so we obtain

$$a^2 + b^2 + 2bt \leq \|T\|^2 \quad \forall t \in \mathbb{R}$$

and so $b = 0$.

□

Lemma 0.59. *Let \mathcal{H} be Hilbert over \mathbb{C} . Let $T \in \mathcal{L}(\mathcal{H})$, and let*

$$M_\lambda = \{x \in \mathcal{H} \mid Tx = \lambda x\} = \text{KER } \lambda I - T$$

be the λ -eigenspace of T . Then

(a) $M_\lambda \perp M_\mu$ if $\lambda \neq \mu$.

(b) If T is normal, each M_λ is T and T^* invariant. That is,

$$T(M_\lambda) \subseteq M_\lambda, \quad T^*(M_\lambda) \subseteq M_\lambda.$$

PROOF.

(a) Let $u \in M_\lambda, v \in M_\mu$. Then

$$\begin{aligned} (\lambda - \mu) \langle u, v \rangle &= \langle \lambda u, v \rangle - \langle u, \bar{\mu} v \rangle \\ &= \langle Tu, v \rangle - \langle u, T^* v \rangle \\ &= \langle Tu, v \rangle - \langle Tu, v \rangle \\ &= 0 \end{aligned}$$

and so $\langle u, v \rangle = 0$.

(b) If T is normal, then $\text{KER } T = \text{KER } T^*$ as

$$\begin{aligned} \|Tx\|^2 &= \langle Tx, Tx \rangle = \langle x, T^*Tx \rangle \\ &= \langle x, TT^*x \rangle = \langle T^*x, T^*x \rangle \\ &= \|T^*x\|^2. \end{aligned}$$

Similarly, if T is normal then $\lambda I - T$ is normal. Then

$$\begin{aligned} M_\lambda &= \text{KER } \lambda I - T \quad (T \text{ invariant}) \\ &= \text{KER } \bar{\lambda} I - T^* \quad (T^* \text{ invariant}). \end{aligned}$$

□

The spectral theory for compact normal operators in a Hilbert space is particularly nice, as the following theorem demonstrates.

Theorem 0.60. *Let $T \in \mathcal{L}(\mathcal{H})$ be compact and normal. Then*

$$\mathcal{H} = \overline{\bigoplus_{\lambda \in \sigma(T)} M_\lambda},$$

the closure of the span of the eigenspaces, and \mathcal{H} has an orthonormal basis consisting of eigenvectors. Moreover, T acts diagonally with respect to this basis.

PROOF. Let

$$M = \overline{\bigoplus_{\lambda \in \sigma(T)} M_\lambda},$$

a closed subspace. Hence $H = M \oplus M^\perp$, where

$$M^\perp = \{x \in \mathcal{H} \mid \langle x, m \rangle = 0 \forall m \in M\}.$$

We must show that $M^\perp = \{0\}$. Assume the contrary. Then consider $\tilde{T} = M^\perp \rightarrow \mathcal{H}$ be the restriction of T to M^\perp . Then we have

$$\tilde{T} : M^\perp \rightarrow M^\perp$$

is compact and normal (Exercise). Then

- (a) $\sigma(\tilde{T}) = \{0\}$. Then $r(\tilde{T}) = 0$, and so $\|\tilde{T}\| = 0$, and so $\tilde{T} = 0$. Then each $x \in M^\perp \setminus \{0\}$ satisfies $\tilde{T}x = 0 = 0x$, and so $x \in M_0$ with $M^\perp \subseteq M_0 \subseteq M$, a contradiction (from direct sum decomposition). Hence $M = \{0\}$.
- (b) $\sigma(\tilde{T}) \neq \{0\}$. So there is an eigenvalue $\lambda \in \sigma(T) \setminus \{0\}$. So there is $x \in M^\perp \setminus \{0\}$ with $\tilde{T}x = \lambda x$. o $Tx = \lambda x$, and so $x \in (M_\lambda \cap M^\perp) \setminus \{0\}$, a contradiction. Hence $M^\perp = \{0\}$.

Choose an orthonormal basis for each M_λ , and combine to get an orthonormal basis of \mathcal{H} , using $M_\lambda \perp M_\mu$. □

1. The Fredholm Alternative

Recall that for matrices, we have the following result, known as the Fredholm alternative.

Theorem 1.1 (Fredholm alternative (Finite dimensional spaces)). *Let $A : \mathbb{C}^n \rightarrow \mathbb{C}^n$ be linear. Then exactly one of the following two things occur:*

- (1) $Ax = 0$ has only the trivial solution $x = 0$, in which case $Ax = b$ has a unique solution for each $b \in \mathbb{C}^n$.
- (2) $Ax = 0$ has a non-trivial solution, in which case $Ax = b$ has either no solutions, or infinitely many solutions.

Definition 1.2 (Hilbert-Schmidt integral operators).

$$T : L^2([a, b]) \rightarrow L^2([a, b])$$

$$(Tf)(x) \mapsto \int_a^b K(x, y)f(y) dy$$

where $\|K\|_2$ is finite. These are compact operators.

Consider equations of the following form

$$\lambda f(x) - \int_a^b K(x, y)f(y) dy = g(x),$$

where $\lambda \neq 0$ and $g \in L^2$ are given. This can be rewritten in the form

$$(\lambda I - T)f = g.$$

Then we have the following theorem, due to Fredholm.

Theorem 1.3 (Fredholm alternative (Hilbert spaces)). *Let \mathcal{H} be Hilbert over \mathbb{C} , and let $T \in \mathcal{K}(\mathcal{H})$. Then exactly one of the following occurs.*

- (a) $(\lambda I - T)x = 0$ has only the trivial solution, in which case $(\lambda I - T)x = b$ has a unique solution for each $b \in \mathcal{H}$.
- (b) $(\lambda I - T)x = 0$ has a non trivial solution, in which case $(\lambda I - T)x = b$ has a solution if and only if $b \perp y$ for every solution y of the equation

$$(\bar{\lambda}I - T^*)y = 0$$

This is finite dimensional, as it is the kernel of $(\lambda I - T)^$.*

PROOF.

- (a) If $(\lambda I - T)x = 0$ has only the trivial solution, then $\text{KER } \lambda I - T = \{0\}$ and so it is injective. Hence λ is not an eigenvalue, and so λ is not a spectral value. So $\lambda I - T$ is invertible, and so $(\lambda I - T)x = b$ has a unique solution $x = (\lambda I - T)^{-1}b$, which can be expanded into a series expression if $|\lambda| > r(T)$.
- (b) Suppose $(\lambda I - T)x = 0$ has a non-trivial solution. Then

$$\begin{aligned} & (\lambda I - T)x = b \text{ has a solution} \\ \iff & b \in \text{IM } \lambda I - T \text{ which is closed} \\ \iff & b \in ((\text{IM } \lambda I - T)^\perp)^\perp \\ \iff & b \in (\text{KER } \bar{\lambda} - T^*)^\perp \\ \iff & b \perp y \quad \forall y \in \text{KER } \bar{\lambda}I - T^*. \quad \square \end{aligned}$$

Proposition 1.4 (Miscellaneous).

- (a) If M is a closed subspace of \mathcal{H} , then $M = M^{\perp\perp}$.
 (b) If $S : \mathcal{H} \rightarrow \mathcal{H}$ and $S \in \mathcal{L}(\mathcal{H})$, then $(\text{IM } S)^{\perp} = \text{KER } S^*$.

PROOF.

- (a) Let $m \in M$, then $\langle m, x \rangle = 0$ for all $x \in M^{\perp}$, and so $m \in (M^{\perp})^{\perp} = M^{\perp\perp}$, and so $M \subseteq M^{\perp\perp}$.
 Let $x \in M^{\perp\perp}$. Since M is closed, $\mathcal{H} = M \oplus M^{\perp}$, and so $x = m + m^{\perp}$. So $x - m \in M^{\perp\perp} + M \subseteq M^{\perp\perp}$, and so $x - m = m^{\perp} \in M^{\perp\perp}$. But M^{\perp} is closed, and so $\mathcal{H} = M^{\perp} \oplus M^{\perp\perp}$. So $x - m = 0$, and $x = m \in M$.
 (b)

$$\begin{aligned}
 (\text{IM } S)^{\perp} &= \{x \in \mathcal{H} \mid \langle x, sy \rangle = 0 \quad \forall y \in \mathcal{H}\} \\
 &= \{x \in \mathcal{H} \mid \langle S^*x, y \rangle = 0 \quad \forall y \in \mathcal{H}\} \\
 &= \{x \in \mathcal{H} \mid S^*x = 0\} \\
 &= \text{KER } S^*
 \end{aligned}$$

□